

PERCENTAGE COMPOSITION

- sometimes called "percent composition" or "percent composition by mass"
- the percentage of each element in a compound, expressed in terms of mass

Example: Find the percentage composition of ammonium nitrate.

$$\text{NH}_4\text{NO}_3 : \text{N} : 2 \times 14.01 = 28.02$$



$$\text{H} : 4 \times 1.008 = 4.032$$

$$\text{O} : 3 \times 16.00 = 48.00$$

$$\overline{80.052 \text{ g NH}_4\text{NO}_3 \text{ per mole}}$$

$$\% \text{ N} = \frac{28.02}{80.052} \times 100\% = 35.0\% \text{ N}$$

$$\% \text{ H} = \frac{4.032}{80.052} \times 100\% = 5.0\% \text{ H}$$

$$\% \text{ O} = \frac{48.00}{80.052} \times 100\% = 60.0\% \text{ O}$$

So far, we have

- looked at how to determine the composition by mass of a compound from a formula
- converted from MASS to MOLES (related to the number of atoms/molecules)
- converted from MOLES to MASS

Are we missing anything?

- What about SOLUTIONS, where the desired chemical is not PURE, but found DISSOLVED IN WATER?
- How do we deal with finding the moles of a desired chemical when it's in solution?

MOLAR CONCENTRATION

- unit: MOLARITY (M): moles of dissolved substance per LITER of solution

$$M = \text{molarity} = \frac{\text{moles of solute}}{\text{L solution}}$$

6.0 M HCl solution: $\frac{6.0 \text{ mol HCl}}{\text{L}}$

If you have 0.250 L (250 mL) of 6.0 M HCl, how many moles of HCl do you have?

$$6.0 \text{ mol HCl} = 1 \text{ L solution}$$

$$0.250 \text{ L solution} \times \frac{6.0 \text{ mol HCl}}{1 \text{ L solution}} = 1.5 \text{ mol HCl}$$

If you need 0.657 moles of hydrochloric acid, how many liters of 0.0555 M HCl do you need to measure out?

$$0.0555 \text{ mol HCl} = 1 \text{ L solution}$$

$$0.657 \text{ mol HCl} \times \frac{1 \text{ L solution}}{0.0555 \text{ mol HCl}} = 11.8 \text{ L solution}$$


What if we used 6.00 M HCl?

$$6.00 \text{ mol HCl} = 1 \text{ L solution}$$



$$0.657 \text{ mol HCl} \times \frac{1 \text{ L solution}}{6.00 \text{ mol HCl}} = 0.110 \text{ L solution} \\ (110 \text{ mL})$$

CHEMICAL CALCULATIONS CONTINUED: REACTIONS

- Chemical reactions proceed on an ATOMIC basis, NOT a mass basis!
- To calculate with chemical reactions (i.e. use chemical equations), we need everything in terms of ATOMS ... which means MOLES of atoms




coefficients are in terms of atoms and molecules!



- To do chemical calculations, we need to:

- Relate the amount of substance we know (mass or volume) to a number of moles
- Relate the moles of one substance to the moles of another using the equation
- Convert the moles of the new substance to mass or volume as desired



* Given that we have 25.0 g of liquid bromine, how many grams of aluminum would we need to react away all of the bromine? How many grams of aluminum bromide would be produced?

(1) Convert grams of bromine to moles: Need formula weight $\text{Br}_2 : \frac{2 \times 79.90}{159.80}$

$$159.80 \text{ g Br}_2 = 1 \text{ mol Br}_2$$

$$25.0 \text{ g Br}_2 \times \frac{1 \text{ mol Br}_2}{159.80 \text{ g Br}_2} = 0.15645 \text{ mol Br}_2$$

(2) Use the chemical equation to relate moles of bromine to moles of aluminum

$$2 \text{ mol Al} = 3 \text{ mol Br}_2$$

$$0.15645 \text{ mol Br}_2 \times \frac{2 \text{ mol Al}}{3 \text{ mol Br}_2} = 0.10430 \text{ mol Al}$$

(3) Convert moles aluminum to mass: Need formula weight $\text{Al} : 26.98$


$$26.98 \text{ g Al} = 1 \text{ mol Al}$$

$$0.10430 \text{ mol Al} \times \frac{26.98 \text{ g Al}}{1 \text{ mol Al}} = \boxed{2.81 \text{ g Al}}$$

You can combine all three steps on one line if you like!

$$25.0 \text{ g Br}_2 \times \frac{1 \text{ mol Br}_2}{159.80 \text{ g Br}_2} \times \frac{2 \text{ mol Al}}{3 \text{ mol Br}_2} \times \frac{26.98 \text{ g Al}}{1 \text{ mol Al}} = 2.81 \text{ g Al}$$

(1) (2) (3)



But ...

... what would you have done to calculate the mass of aluminum bromide IF you had NOT been asked to calculate the mass of aluminum FIRST?


$$25.0 \text{ g Br}_2 \times \frac{1 \text{ mol Br}_2}{159.80 \text{ g Br}_2} \times \frac{2 \text{ mol AlBr}_3}{3 \text{ mol Br}_2} \times \frac{266.68 \text{ g AlBr}_3}{1 \text{ mol AlBr}_3} = 27.8 \text{ g AlBr}_3$$

$$\text{AlBr}_3 : \text{Al} : 1 \times 26.98$$

$$\begin{array}{r} \text{Br} : 3 \times 79.90 \\ \hline 266.68 \end{array}$$

Example:

How many milliliters of 6.00M hydrochloric acid is needed to completely react with 25.0 g of sodium carbonate?



- Convert mass of sodium carbonate to moles using formula weight
 - Convert moles of sodium carbonate to moles hydrochloric acid using chemical equation
 - Convert moles of hydrochloric acid to volume using concentration (M = moles/L)
-

- Convert mass of sodium carbonate to moles using formula weight


$$\begin{aligned}\text{Na}_2\text{CO}_3: \quad \text{Na} &: 2 \times 22.99 \\ &\text{C} : 1 \times 12.01 \\ &\text{O} : 3 \times 16.00 \\ \hline &105.99\end{aligned}$$

$$105.99 \text{ g Na}_2\text{CO}_3 = 1 \text{ mol Na}_2\text{CO}_3$$

$$25.0 \text{ g Na}_2\text{CO}_3 \times \frac{1 \text{ mol Na}_2\text{CO}_3}{105.99 \text{ g Na}_2\text{CO}_3} = 0.23587 \text{ mol Na}_2\text{CO}_3$$

Example:

How many milliliters of 6.00M hydrochloric acid is needed to completely react with 25.0 g of sodium carbonate?



- Convert moles of sodium carbonate to moles hydrochloric acid using chemical equation

$$2 \text{ mol HCl} = 1 \text{ mol Na}_2\text{CO}_3$$
$$0.23587 \frac{1 \text{ mol Na}_2\text{CO}_3}{1 \text{ mol Na}_2\text{CO}_3} \times \frac{2 \text{ mol HCl}}{1 \text{ mol Na}_2\text{CO}_3} = 0.471743 \text{ mol HCl}$$

- Convert moles of hydrochloric acid to volume using concentration (M = moles/L)

$$6.00 \text{ M HCl}: \quad 6.00 \text{ mol HCl} = 1 \text{ L}$$


$$0.471743 \text{ mol HCl} \times \frac{1 \text{ L}}{6.00 \text{ mol HCl}} = 0.0786 \text{ L of solution}$$

$$\text{mL} = 10^{-3} \text{ L} \quad \text{Convert liters to milliliters!}$$

$$0.0786 \text{ L of solution} \times \frac{\text{mL}}{10^{-3} \text{ L}} = 78.6 \text{ mL solution}$$

CONCEPT OF LIMITING REACTANT

- When does a chemical reaction STOP?



- When does this reaction stop? When burned in open air, this reaction stops when all the MAGNESIUM STRIP is gone. We say that the magnesium is LIMITING.
- This reaction is controlled by the amount of available magnesium
- At the end of a chemical reaction, the LIMITING REACTANT will be completely consumed, but there may be amount of OTHER reactants remaining. We do chemical calculations in part to minimize these "leftovers".