
More on precison

Every measurement contains some amount of ERROR, or some amount of deviation from the true value of what is being measured.

RANDOM ERROR is the variability in a measurement that cannot be traced back to a single cause. Random errors cause measurements to fluctuate around the true value, but can be averaged out given enough measurements.

When reporting measurements, we want to indicate how much random error we think is present. How?



How long is the green line?

For this experiment, measure the line and record your answer in the form: X.XX cm (In other words, measure to the nearest 0.01 cm)

Write your answer on the sheet, then fold the sheet in half and pass it up to the front!

After throwing away obvious mistakes in reading the scale, we had:

Value	# students
1,60	2
1.61	2
1.62	2
1,63	8
1.64	

1,62266667 cm: unrounded average Overall average $1.62 \pm .01$ cm

Certain. Unce Little to no Expe variation vary expected. +/- 1 Same almost every time

Uncertain. Expected to vary by about +/- 1

15 measurements

When reading measurements from a scale, record all CERTAIN digits and one UNCERTAIN (or estimated) digit.

Significant figures

SIGNIFICANT FIGURES are a way to indicate the amount of uncertainty in a measurement.

The significant figures in a measurement are all of the CERTAIN DIGITS plus one and only one UNCERTAIN (or estimated) DIGIT

Determining significant figures

When you read a measurement that someone has written using the significant figures convention, you can tell how precisely that measurement was made.

$$\frac{473}{2} = \pm 0.001g$$
This was measured to the nearest +/-0.001g
The last digit is always UNCERTAIN (or estimated)

$$2 = 1 = \frac{50me \ other \ examples}{3.2076} = \frac{10.0001g}{1 \ uncertain.'}$$


$$37.26 \ kg \pm 0.01 \ kg} = \frac{27.3m}{1 \ uncertain.'}$$

<u>A small problem</u>

The number ZERO has several uses. It may be a measured number, but it may also be a mere "placeholder" that wasn't measured at all!

So how do we tell a measured zero from a placeholder? There are a few ways:

1: BEGINNING ZEROS: Beginning zeros are NEVER considered significant.



2: END ZEROS are sometimes considered significant. They are significant if

- there is a WRITTEN decimal point in the number

10.01 Km

- there is another written indicator that the zero is

significant. Usually this is a line drawn over or under the last zero that is significant!

This zero IS considered significant. There's a written decimal.

± 100 m

These zeros ARE NOT considered significant (no written decimal, and no other indication that the zeros are significant)


M

Σ,

These zeros are not significant.

This zero IS significant. It's marked.

How many significant figures are there in each of these measurements?




(Number of significant figures is indicated in RED below each measurement. Significant digits are UNDERLINED.)

(Approximate uncertainty in each of these measurements is indicated in GREEN after each one.)

Calculations with measurements

When you calculate something using measured numbers., you should try to make sure the ANSWER reflects the quality of the data used to make the calculation.

An ANSWER is only as good as the POOREST measurement that went into finding that answer!




✓ If you add an uncertain number to either a certain or an uncertain number, then the result is uncertain!

 \star If you add certain numbers together, the result is certain!

For addition and subtraction, round FINAL ANSWERS to the same number of decimal places as the measurement with the fewest decimal places. This will give an answer that indicates the proper amount of uncertainty.

For multiplication and division, round FINAL ANSWERS to the same number of SIGNIFICANT FIGURES as the measurement with the fewest SIGNIFICANT FIGURES!



A few more math with significant figures examples:

Placeholder zeros, even though they aren't SIGNIFICANT, still need to be included, so we know how big the number is!

147.<u>3</u> 243<u>2</u> 0.97 + 111.<u>6</u> 2.691.§7 2.692 DENSITY CALCULATION 6 14.7068g 2.7 mL = 5.446962963 ⁹/mL

9

5.4

To improve (make more precise) this calculated density, we must improve the poorest measturement. We must use a more precise device to measure the VOLUME (which only has two significant figures in this example)!

Exact Numbers

- Some numbers do not have any uncertainty. In other words, they weren't measured!

1) Numbers that were determined by COUNTING!


1 H exactly 4 ! 2) Numbers that arise from DEFINITIONS, often involving relationships between units 12 in = 1 ft Km = 10³m & All metric prefixes are exact.

How many blocks are to the left?

- Treat exact numbers as if they have INFINITE significant figures!

Example

You'll need to round the answer to the right number of significant figures!



*An inch is defined as EXACTLY 2.54 cm !

When merely converting the units of a measurement, you almost always have the same number of significant figures in the answer as you did in the original measurement. (EXCEPTION: Temperature conversions, since they involve addition and subtraction)