



A DOT STRUCTURE FOR A POLYATOMIC ION

- 1) Count valence electrons
- 2) Pick central atom and draw skeletal structure
 - central atom is usually the one that needs to gain the most electrons!
 - skeletal structure has all atoms connected to center with single bonds
- 3) Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.
- 4) Check octet rule - each atom should have a share in 8 electrons (H gets 2). if not, make double or triple bonds.


$$\underline{\quad 9 \quad}$$
$$- 1$$
$$\underline{\quad 8 \quad}$$

An odd number of electrons? But Lewis structures deal in PAIRS of electrons!


Subtract an electron to account for ammonium's +1 charge



Draw brackets around the structure so that we can see the entire molecule is CHARGED.

EXPANDED VALENCE and other exceptions to the "octet rule"

- Some atoms do not always obey the octet rule. A few, like BORON, will bond in such a way that they end up with less than eight electrons.




... but many more bond in such a way that they end up with a share in MORE THAN EIGHT electrons!

- Any atom in period three or greater can do this. SULFUR and PHOSPHORUS compounds commonly do this!

... these atoms have unfilled "d" orbitals that may participate in bonding!

- All noble gas compounds (example: XENON compounds with oxygen and fluorine) exhibit this behavior!

EXAMPLES:




$$\begin{array}{l} \text{S: } 6 \\ \text{F: } \frac{7 \times 6}{48} \end{array}$$

- The central SULFUR atom has a share in TWELVE total electrons, not eight!
- The SHAPE of the sulfur hexafluoride molecule in three dimensions agrees with the picture of six fluorine atoms each sharing a pair of electrons with a sulfur center.




$$\begin{array}{l} \text{P: } 5 \\ \text{Cl: } \frac{7 \times 3 = 21}{26} \end{array}$$



This structure obeys the octet rule.



$$\begin{array}{l} \text{P: } 5 \\ \text{Cl: } \frac{7 \times 5 = 35}{40} \end{array}$$



This molecule does NOT obey the octet rule. Phosphorus ends up with ten electrons instead of eight.

FORMAL CHARGE

- You can often draw more than one structure for a molecule that appears correct. How can you determine which one is more likely?

- USE FORMAL CHARGE!

- Formal charge is a hypothetical charge on each atom in a structure. It assumes:

- ① All bonding electrons are shared EQUALLY between atoms
- ② Lone pairs are NOT shared.


$$\text{FORMAL CHARGE} = \text{ORIGINAL \# OF VALENCE ELECTRONS} - \text{NUMBER OF BONDS} - \text{NUMBER OF UNSHARED ELECTRONS}$$

* The sum of the formal charges of all atoms in a structure should equal to the charge of the molecule (0 for neutral molecules)

The "better" Lewis structure will have:

- Lower magnitudes of formal charge (0 0 is better than +2 -2)
- Negative formal charges on ELECTRONEGATIVE atoms, or positive formal charges on atoms that are less electronegative.

EXAMPLE: COCl_2



... calculate formal charges to tell which structure is more likely!

$$\text{O: } 6 - 2 - 4 = 0$$

$$\text{C: } 4 - 4 - 0 = 0$$

$$\text{Cl: } 7 - 1 - 6 = 0$$

$$\text{Cl: } 7 - 1 - 6 = 0$$

$$\text{O: } 6 - 1 - 6 = -1$$

$$\text{C: } 4 - 4 - 0 = 0$$

$$=\text{Cl: } 7 - 2 - 4 = +1$$

$$-\text{Cl: } 7 - 1 - 6 = 0$$

* The sum of the formal charges on both structures equals zero - so we've drawn them correctly. (This is a neutral molecule, not a polyatomic ion)

* The structure on the left is preferred. It has LOWER formal charges (all zeros) than the structure on the right (+1/-1/0). The structure on the right also has a +1 formal charge on electronegative CHLORINE.



... we can determine which of these structures is more likely by calculating formal charges!

$$\text{H}: 1 - 1 - 0 = 0$$

$$\text{C}: 4 - 3 - 2 = -1$$

$$\text{N}: 5 - 4 - 0 = +1$$

$$\text{H}: 1 - 1 - 0 = 0$$

$$\text{C}: 4 - 4 - 0 = 0$$

$$\text{N}: 5 - 3 - 2 = 0$$

Which structure is more likely?

- * The HCN structure (right) is more likely. It has lower formal charges (all zeros).
- Also, the structure on the left gives a positive formal charge to electronegative NITROGEN (while the less electronegative CARBON gets a negative formal charge).

Let's look at sulfur trioxide.







$$\text{S} : 6$$


$$\text{O} : 6 \times 3 = 18$$

$$\hline 24 e^-$$

Skeletal structure:




Resonance structures.



Expanded valence
(Sulfur is period 3)

To decide which structure is preferred, let's look at formal charges.




Resonance structures.

$$S: 6 - 4 - 0 = +2$$

$$O - : 6 - 1 - 6 = -1$$

$$O - : 6 - 1 - 6 = -1$$

$$O = : 6 - 2 - 4 = 0$$



Expanded valence
(Sulfur is period 3)

$$S: 6 - 6 - 0 = 0$$

$$O = : 6 - 2 - 4 = 0$$

$$O = : 6 - 2 - 4 = 0$$

$$O = : 6 - 2 - 4 = 0$$

BASED ON FORMAL CHARGES, the expanded valence structure is preferred.

The correct structure is typically the one with minimized formal charges - even if it violates the octet rule. (Just remember - period 2 NEVER violates this octet rule - no unfilled "d" orbitals)