

$$l_2 SO_4(a_q) \rightarrow NO REACTION$$


Silver is LESS ACTIVE than hydrogen, so we expect that no reaction will occur. Silver is not able to give electrons to hydrogen.

$$M_g(s) + Z_n SOy(u_q) \rightarrow M_g SOy(u_q) + Z_n(s)$$

Magnesium is MORE ACTIVE than zinc, so we would expect a replacement reaction to occur.

179 (5)+

CHEMICAL CALCULATIONS - RELATING MASS AND ATOMS



- While chemical equations are written in terms of ATOMS and MOLECULES, that's NOT how we often measure substances in lab!


- measurements are usually MASS (and sometimes VOLUME), NOT number of atoms or molecules!  $\bigwedge_{Na2} Co_3 solid$   $\bigwedge_{Hcl} Solution$ 

... so how do we relate atoms and molecules with things we routinely measure in lab - like grams and milliliters?

## THE MOLE CONCEPT

- A "mole" of atoms is  $6.022 \times 10^{23}$  why so big? Because atoms are so small!


- Why - in the metric dominated world of science - do we use such a strange number for quantity of atoms?



## THE MOLE CONCEPT

- Why define the mole based on an experimentally-measured number?

- The atomic weight of an element (if you put the number in front of the unit GRAMS) is equal to the mass of ONE MOLE of atoms of that element!



Magnesium (Mg): 24.31 g = the mass of ONE MOLE OF MAGNESIUM ATOMS

- So, using the MOLE, we can directly relate a mass and a certain number of atoms!

RELATING MASS AND MOLES

- Use DIMENSIONAL ANALYSIS (a.k.a "drag and drop")
- Need CONVERSION FACTORS where do they come from?
- We use ATOMIC WEIGHT as a conversion factor.

$$M_{g} : 2H.31 \qquad 2H.31 g M_{g} = 1 \mod M_{g}$$
  
"mol" is the abbreviation for "mole"  
nple: How many moles of atoms are there in 250. g of magnesium metal?  
2H.31 g Mg = mol Mg  
2S0. g Mg x  $\frac{mol Mg}{2H.31 g Mg} = 10.3 \mod Mg$ 

Example: