

Bohr's model didn't account for electron-electron interactions (which didn't exist in HYDROGEN)


- To account for this added complexity, a more sophisticated model had to be devised: QUANTUM THEORY. Quantum theory is the modern picture of the atom and its electron cloud.

SHELLS, SUBSHELLS, AND ORBITALS

- Bohr's model predicted that energy levels (called SHELLS) were enough to describe completely how electrons were arranged around an atom. But there's more to it!

SHELL: Equivalent to Bohr's energy levels. Electrons in the same SHELL are all the same distance from the nucleus. They all have SIMILAR (but not necessarily the SAME) energy.


- Shells are numbered (1-... - Elements on the periodic table have shells numbered from 1 to 7)
- Higher numbers correspond to greater distance from the nucleus and greater energy, and larger size!
- Higher shells can hold more electrons than lower shells!




SUBSHELLS: Within a SHELL, electrons may move in different ways around the nucleus! These different "paths" are called SUBSHELLS

- SHAPES of regions of space that electrons are able to exist in.


Illustrations:
p315-316



"s" subshell
(a spherical region)



"p" subshell
(a dumbbell shaped region)



"d" subshell

- Some atoms also have "f" subshells (not pictured)

See p 314-316 for nicer drawings of the subshells.

ORBITALS - are specific regions of space where electrons may exist

- The SHAPE of an orbital is defined by the SUBSHELL it is in
- The ENERGY of an orbital is defined by both the SHELL the orbital is in AND the kind of SUBSHELL it is in
- Each orbital may, at most, contain TWO ELECTRONS

ARRANGEMENT OF SHELLS, SUBSHELLS, AND ORBITALS

- Shells are numbered. Each shell can contain the same number of SUBSHELLS as its number:

1st shell: ONE possible subshell (s)


2nd shell: TWO possible subshells (s, p)

3rd shell: THREE possible subshells (s, p, d)

4th shell: FOUR possible subshells (s, p, d, f)




... and so on

- Each subshell can contain one or more ORBITALS, depending on how many different ways there are to arrange an orbital of that shape around the nucleus.



"s" subshell
One possible
orientation

Maximum 2 electrons in 1 orbital



"p" subshell: Three possible orientations

Maximum 6 electrons in 3 orbitals


- There are five possible orbitals in a "d" subshell, and 7 possible orbitals in an "f" subshell!

Maximum 10 electrons
in 5 orbitals
(see p 316)

Maximum 14 electrons
in 7 orbitals

ENERGY DIAGRAM

- We can map out electrons around an atom using an energy diagram:




Let's look at some example atoms:


Magnesium: $Z=12$, 12 electrons

$\uparrow Z$: atomic #

ENERGY




Outermost electrons of magnesium "valence electrons". These electrons are involved in chemical bonding!

Aluminum: $Z = 13$ 

Aluminum has THREE valence electrons!
(All electrons in the outer shell are valence electrons!)




Atoms tend to form ions or chemical bonds in order to end up with filled outer "s" and "p" subshells.

This is called the "octet" rule. (Not all chemical bonds follow this - it's a RULE OF THUMB, not a scientific law!)

Example: Oxygen, $Z = 8$ 

ELECTRON CONFIGURATION

- A shorthand way to write about electron arrangement around an atom.



↑ Valence electrons are the ones in the outermost SHELL, not just the last subshell. Aluminum has THREE valence electrons.

two
elements
wide
IA

ELECTRON CONFIGURATION AND THE PERIODIC TABLE

Helium is part
of the "s" block!

1	H	IIA											six elements wide						He
2	Li	Be											B	C	N	O	F	Ne	
3	Na	Mg	ten elements wide										Al	Si	P	S	Cl	Ar	
4	K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr	
5	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe	
6	Cs	Ba	La*	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn	
7	Fr	Ra	Ac*	Rf	Db	Sg	Bh	Hs	Mt	*inner transition metals go here									

"s" block: last electron in these atoms is in an "s" orbital!

"p" block: last electron in these atoms is in a "p" orbital!


"d" block: last electron in these atoms is in a "d" orbital

- To write an electron configuration using the periodic table, start at hydrogen, and count up the electrons until you reach your element!



1	IA H											VIII A He						
2	II A Li	Be											III A B	IV A C	V A N	VII A O	VII A F	Ne
3	III B Na	Mg	III B	IV B	V B	VII B	VII B	VIII B	IB	IIB	Al	Si	P	S	Cl	Ar		
4	K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
5	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
6	Cs	Ba	La*	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
7	Fr	Ra	Ac*	Rf	Db	Sg	Bh	Hs	Mt	*"inner" transition metals go here								

Example: Phosphorus (P): $1s^2 2s^2 2p^6 3s^2 3p^3$


EXAMPLES:




Remember - valence electrons are ALL of the electrons in the outermost SHELL! (may have more than one SUBSHELL)!



TITANIUM is a transition metal that commonly forms either +2 or +4 cations. The 4s electrons are lost when the +2 ion forms, while the 4s AND 3d electrons are lost to form the +4!



Noble gas core notation. Use the previous noble gas on the table, then add the electrons that it doesn't have to the end.




You are responsible for writing electron configurations up to Z=18, Argon. These are here to illustrate other points!

PERIODIC TRENDS

- Some properties of elements can be related to their positions on the periodic table.

ATOMIC RADIUS


- The distance between the nucleus of the atoms and the outermost shell of the electron cloud.
- Relates to the size of the atom.
- As you go DOWN A GROUP (↓), the atomic radius INCREASES.



- Why? As you go down a period, you are ADDING SHELLS!


- As you go ACROSS A PERIOD (→), the atomic radius DECREASES

Why? Let's look at some sample atoms.



lithium

Outer electron sees an effective +1 charge (shielded from +3 nucleus by 2 electrons)



fluorine

Outer electrons see an effective +7 charge (shielded from +9 nucleus by 2 electrons)

... so fluorine's outer shell is pulled closer to the nucleus than lithium's!



IONIZATION ENERGY (or FIRST IONIZATION ENERGY)

- The amount of energy required to remove a single electron from the outer shell of an atom.
- Relates to reactivity for metals. The easier it is to remove an electron, the more reactive the metal.
- As you go DOWN A GROUP (\downarrow), the ionization energy DECREASES.

- Why? As you go down a period, you are ADDING SHELLS. Since the outer electrons are farther from the nucleus and charge attraction lessens with distance, this makes electrons easier to remove as the atoms get bigger!

- As you go ACROSS A PERIOD (\longrightarrow), the ionization energy INCREASES.

- Why? Let's look at some sample atoms.



... since fluorine's outer electrons are held on by a larger effective charge, they are more difficult to remove than lithium's.

PERIODIC TRENDS IN A NUTSHELL

LARGER
IONIZATION
ENERGY

SMALLER
RADIUS

IA												VIII A					
H	IIA											III A	IVA	VA	VIA	VIIA	He
Li	Be											B	C	N	O	F	Ne
Na	Mg	IIIB	IVB	VB	VIB	VII B	VIII B		IB	IIB	Al	Si	P	S	Cl	Ar	
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
Cs	Ba	La*	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
Fr	Ra	Ac*	Rf	Db	Sg	Bh	Hs	Mt	*"inner" transition metals go here								

LARGER
RADIUS

SMALLER
IONIZATION
ENERGY


... but using electron configurations to describe how aluminum bromide forms is a bit cumbersome! Can we simplify the picture a bit?

LEWIS NOTATION / ELECTRON-DOT NOTATION

- Lewis notation represents each VALENCE electron with a DOT drawn around the atomic symbol. Since the valence shell of an atom contains only "s" and "p" electrons, the maximum number of dots drawn will be EIGHT.

- To use electron-dot notation, put a dot for each valence electron around the atomic symbol. Put one dot on each "side" of the symbol (4 sides), then pair the dots for atoms that have more than four valence electrons.

examples:



Which "side" you draw the dots on isn't important, as long as you have the right number of electrons and the right number of "pairs"



... are all equivalent!

To draw a dot structure for an atom, you need to know HOW MANY valence electrons it has! You can determine this simply from the periodic table, WITHOUT writing the whole electron configuration!

The number of valence electrons equals the group number in the A/B group numbering system

	IA																	VIIIA
1	H	IIA										IIIA	IVA	VA	VIA	VIIA		He
2	Li	Be										B	C	N	O	F		Ne
3	Na	Mg	IIIB	IVB	VB	VIB	VII B	VIII B	IB	IIB		Al	Si	P	S	Cl		Ar
4	K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
5	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
6	Cs	Ba	La*	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
7	Fr	Ra	Ac*	Rf	Db	Sg	Bh	Hs	Mt									

1 valence electron

2 valence electrons

3 valence electrons

4 valence electrons


5 valence electrons

6 valence electrons

7 valence electrons

8 valence electrons (except helium!)

... but how do we use this to describe a reaction that produces ions? Let's look at our previous example!




... this is a bit easier to follow than looking at all those letters and numbers in the electron configurations for these elements!


MOLECULAR COMPOUNDS

- Form when atoms SHARE electrons instead of transferring them. This results in the formation of MOLECULES ... groups of atoms held together by electron-sharing.

How might atoms SHARE electrons? By coming together close enough so that their atomic ORBITALS overlap each other:



Each hydrogen atom has a single electron in a 1s orbital.



When hydrogen atoms come close enough to each other for these orbitals to OVERLAP, each hydrogen "sees" BOTH electrons, filling up the "s" orbitals of both atoms. This is a COVALENT BOND.

— This is the DIATOMIC MOLECULE, H_2