
$\widehat{i}$  Atoms may share one, two, or three pairs of electrons.

Atoms will usually share enough electrons so that each atom ends up with a share in EIGHT electrons - the "octet rule"

- HYDROGEN will only end up with two electrons!

- Some other atoms may end up with more or less than eight electrons. Exceptions to the octet rule are covered in Chapter 9.

NOW, how could we come up with dot structures for some more complicated (and therefore, more interesting) molecules?



<sup>205</sup> DRAWING DOT STRUCTURES FOR SIMPLE MOLECULES

) Count valence electrons

Pick central atom and draw skeletal structure

central atom is usually the one that needs to gain the most electrons!
skeletal structure

has all atoms connected to center with single bonds

3 Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.


Check octet rule - each atom should have a share in 8 electrons (H gets 2). if not, make double or triple bonds.  $OU_2$ 

$$O: 6$$

$$C[: \frac{7 \times 2}{2}]$$

c: 4

i - c - c



Choose carbon as the central atom and draw skeletal structure.

Distribute remaining electrons. We stop here because we've used all 24 electrons.

- Carbon has only SIX electrons!

\_ We'll pick OXYGEN to share two pairs of electrons, since it needed an additional two electrons from the beginning.

This structure looks better. All atoms have a share in the correct number of electrons!



Dick central atom and draw skeletal structure

- central atom is usually the one that needs to gain the most electrons!

- skeletal structure has all atoms connected to center with single bonds

) Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.

Check octet rule - each atom should have a share in 8 electrons (H gets 2). if not, make double or triple bonds.  $N: S \times I$   $0: 6 \times I$   $C_1: 7 \times I$  $IGe^{-1}$ 

NOCI

○ - ↓ - C ↓
We'll use NITROGEN for our central atom, since it needs to gain 3 more electrons - more than either oxygen or chlorine.

We ran out of space on the outer atoms, so we put the last pair of electrons on the central nitrogen.

NITROGEN has a share in only six electrons in this structure, so we need to make a double bond.



We use a pair of electrons from OXYGEN (same reason as the previous example)

3


## Count valence electrons

Pick central atom and draw skeletal structure

> - central atom is usually the one that needs to gain the most electrons! - skeletal structure has all atoms connected to center with single bonds

Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.

Check octet rule - each atom should have a share in 8 electrons (H gets 2). if not, make double or triple bonds.



These two oxygen atoms SHOULD bond the  $:O \equiv C = 0$ ; same way to the carbon center. They are identical atoms in an identical environment.

EXPERIMNETALLY, we find that the two oxygen atoms are the same distance from he carbon center, so they should both have the same kind of bond with the carbon. Count valence electrons

(2) P


Pick central atom and draw skeletal structure

central atom is usually the one that needs to gain the most electrons!
skeletal structure has all atoms connected to center with single bonds

3 Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.

Ч Check octet rule - each atom should have a share in 8 electrons (H gets 2). if not, make double or triple bonds.

HNO, "nitrous acid" - In oxyacids, the acidic hydrogen atoms are attached to OXYGEN atoms in the structure! H:|X|18e-N'SXI 0:682 0 - N - 0 - H-H ... but NITROGEN has a share in only SIX electrons! :0 = N - 0 - H



Check octet rule - each atom should have a share in 8 electrons (H gets 2). if not, make double or triple bonds.

A DOT STRUCTURE FOR A MOLECULE WITH DELOCALIZED BONDS

) Count valence electrons

Pick central atom and draw skeletal structure

- central atom is usually the one that needs to gain the most electrons!

- skeletal structure has all atoms connected to center with single bonds

Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.

Check octet rule - each atom should have a share in 8 electrons (H gets 2). if not, make double or triple bonds. 0:386218

 $O_{1}^{2} - O_{1}^{2} - O_{1}^{2}$ ; OUT OF ELECTRONS

(OZONE)

- CEntral oxygen has only six electrons

O = O - O; All atoms have a share in eight electrons!

The structure we drew implies that one of the outer oxygen atoms is closer to the central oxygen atom than the other one.

Experimentally, though, we find the two oxygen atoms to be the SAME distance from the center.

In the ozone molecule, electrons are actually being shared between ALL THREE oxygen atoms at the same time. This is called a DELOCALIZED BOND.



The structures in the green box are called RESONANCE STRUCTURES. The "real" structure of ozone is an "average" of the two resonance structures. The "double bond" electrons in these structures are actually shared between all three oxygen atoms