



LIMITING REACTANT CALCULATIONS

- To find the limiting reactant, calculate how much product would be produced from ALL given reactants. Whichever produces the SMALLEST amount of product is the limiting reactant, and the smallest amount of product is the actual amount of product produced.



If you start with 100. g of each reactant, how much calcium carbide would be produced?



$$100. \text{ g CaO} \times \frac{\text{mol CaO}}{56.08 \text{ g CaO}} \times \frac{\text{mol CaC}_2}{\text{mol CaO}} \times \frac{64.10 \text{ g CaC}_2}{\text{mol CaC}_2} = 114 \text{ g CaC}_2$$



$$100. \text{ g C} \times \frac{\text{mol C}}{12.01 \text{ g C}} \times \frac{\text{mol CaC}_2}{3 \text{ mol C}} \times \frac{64.10 \text{ g CaC}_2}{\text{mol CaC}_2} = 178 \text{ g CaC}_2$$

114 g of calcium carbide product will be produced. (Calcium oxide is limiting, and it runs out when 114 g of product are formed, so the reaction must stop there.)

We say that calcium oxide is "limiting" and carbon is present "in excess".

PERCENT YIELD

- Chemical reactions do not always go to completion! Things may happen that prevent the conversion of reactants to the desired/expected product!


① SIDE REACTIONS:



... so in a low-oxygen environment, you may produce less carbon dioxide than expected!

② TRANSFER AND OTHER LOSSES

- When isolating a product, losses may occur in the process. Example: filtering



③ EQUILIBRIUM

- Reactions may reach an equilibrium between products and reactants. We'll talk more about this in CHM 111. The net result is that the reaction will appear to stop before all reactants have been consumed!


- All of these factors cause a chemical reaction to produce LESS product than calculated. For many reactions, this difference isn't significant. But for others, we need to report the PERCENT YIELD.

$$\text{PERCENT YIELD} = \frac{\text{ACTUAL YIELD}}{\text{THEORETICAL YIELD}} \times 100\%$$

↙ Determined EXPERIMENTALLY!


↑ Calculated based on the limiting reactant. (The chemical calculations you've done up to now have been theoretical yields!)

... the percent yield of a reaction can never be greater than 100% due to conservation of mass! If you determine that a percent yield is greater than 100%, then you've made a mistake somewhere - either in a calculation or in the experiment itself!



22.4 grams of benzene are reacted with excess nitric acid. If 31.6 grams of nitrobenzene are collected from the reaction, what is the percent yield?

To find PERCENT YIELD, we need to calculate the THEORETICAL YIELD from the 22.4 grams of benzene we used as a starting material. We already know the actual yield of nitrobenzene.




$$22.4 \text{ g C}_6\text{H}_6 \times \frac{\text{mol C}_6\text{H}_6}{78.114 \text{ g C}_6\text{H}_6} \times \frac{\text{mol C}_6\text{H}_5\text{NO}_2}{\text{mol C}_6\text{H}_6} \times \frac{123.111 \text{ g C}_6\text{H}_5\text{NO}_2}{\text{mol C}_6\text{H}_5\text{NO}_2} = 35.3 \text{ g C}_6\text{H}_5\text{NO}_2$$

\ THEORETICAL YIELD /

$$\% \text{ yield} = \frac{31.6 \text{ g}}{35.3 \text{ g}} \times 100 \% = \boxed{89.5 \%}$$

25.0 mL of acetic acid solution requires 37.3 mL of 0.150 M sodium hydroxide for complete reaction. The equation for this reaction is:



What is the molar concentration of the acetic acid?

$$\frac{\text{mol HC}_2\text{H}_3\text{O}_2}{\text{L Solution}} \leftarrow = 25.0\text{mL or } 0.0250\text{L}$$

Since we already know the VOLUME of the acetic acid, we need to find out how many moles we have to determine the concentration.

$$0.150 \text{ mol NaOH} = \text{L} \quad \text{mol NaOH} = \text{mol HC}_2\text{H}_3\text{O}_2$$


$$0.0373 \text{ L} \times \frac{0.150 \text{ mol NaOH}}{\text{L}} \times \frac{\text{mol HC}_2\text{H}_3\text{O}_2}{\text{mol NaOH}} = 0.005595 \text{ mol HC}_2\text{H}_3\text{O}_2$$

$$\text{Molarity: } \frac{\text{mol HC}_2\text{H}_3\text{O}_2}{\text{L Solution}} = \frac{0.005595 \text{ mol HC}_2\text{H}_3\text{O}_2}{0.0250\text{L}} = \boxed{0.224 \text{ M HC}_2\text{H}_3\text{O}_2}$$

Shortcut: use MILLIMOLES and MILLILITERS instead of moles and liters


$$37.3 \text{ mL} \times \frac{0.150 \text{ mol NaOH}}{\text{L}} \times \frac{\text{mol HC}_2\text{H}_3\text{O}_2}{\text{mol NaOH}} = 5.595 \text{ mmol HC}_2\text{H}_3\text{O}_2$$

$$M = \frac{\text{mol HC}_2\text{H}_3\text{O}_2}{\text{L Solution}} = \frac{5.595 \text{ mmol HC}_2\text{H}_3\text{O}_2}{25.0\text{mL}} = \boxed{0.224 \text{ M HC}_2\text{H}_3\text{O}_2}$$



Calculate how many grams of acrylonitrile could be obtained from 651 kg of propylene, assuming there is excess NO present. (651000 g)

- 1- Convert mass propylene to moles propylene. Use formula weight of propylene
- 2 - Convert moles propylene to moles acrylonitrile. Use coefficients from chemical equation
- 3 - Convert moles acrylonitrile to mass acrylonitrile. Use formula weight of acrylonitrile.



$$651000 \text{ g C}_3\text{H}_6 \times \frac{\text{mol C}_3\text{H}_6}{42,081 \text{ g C}_3\text{H}_6} \times \frac{4 \text{ mol C}_3\text{H}_3\text{N}}{4 \text{ mol C}_3\text{H}_6} \times \frac{53.064 \text{ g C}_3\text{H}_3\text{N}}{\text{mol C}_3\text{H}_3\text{N}} =$$

$$= \boxed{821000 \text{ g C}_3\text{H}_3\text{N}} \quad (821 \text{ kg})$$