
SINGLE REPLACEMENT REACTIONS



... but just because you combine an element and a compound doesn't mean that a reaction will occur. Some combinations react, some don't!

- Whether a reaction occurs depends on how easily the replacing and replaced elements lose electrons. An atom that loses electrons more easily will end up in IONIC form (in other words, in the compound). An atom that loses electrons less easily will end up as a free element.

- We say that an atom that loses electrons more easily that another is MORE ACTIVE than the other element. But how would you get information about ACTIVITY?

A single replacement reaction is an example of a reaction where ELECTRON TRANSFER is a driving force. Electron transfer reactions are generally called OXIDATION-REDUCTION reactions.

ACTIVITY SERIES

- comes from experiental data. It's a list of elements in order of their ACTIVITY - more active elements are higher in the series!

A sample activity series

Sodium
$$Na^{+}$$

Magnesium M_{g}^{2+}
Aluminum $A|^{3+}$
 $Zinc 2n^{3+}$
 $Iron Fe^{2+}$
Lead Pb^{2+}
Hydrogen H^{+}
Copper Cu^{2+}
Silver A_{g}^{+}
Gold Au^{3+}
 $Very active metals will replace
hydrogen in acids AND in
water!
Metals more active than hydrogen
will replace hydrogen in acids!
These metals are
unreactive to most acids!$

PREDICTING SINGLE REPLACEMENT REACTIONS

$$\frac{PREDICTING SINGLE REPLACEMENT REACTIONS}{Plb(s) + 2H(l(nq)) \rightarrow PbC(z(s) + H_2(g))}$$

$$\frac{Pb(s) + 2H(l(nq)) \rightarrow PbC(z(s) + H_2(g))}{Lead is MORE ACTIVE than hydrogen, so we would expect lead to replace hydrogen in this reaction.}$$

$$\frac{Pb(wo_3)_2(nq) + 2n(s) \rightarrow 2n(wo_3)_2(nq) + Pb(s)}{Zinc is MORE ACTIVE than lead, so we expect zinc to replace lead.}$$

$$\frac{Pb(wo_3)_2(nq) + 2n(s) \rightarrow 2n(wo_3)_2(nq) + Pb(s)}{Silver + st}$$

Nat

 $Z_{n^{1+}}$

Fe²⁺

P62+

 H^+

Agt

 Ru^{3+}

A13+

$$A_{\mathcal{G}}(s) + H_{\mathcal{T}}SO_{\mathcal{G}}(\alpha_{\mathcal{G}}) \rightarrow NO REACTION$$


Silver is LESS ACTIVE than hydrogen. Since hydrogen is already in the compound, we do not expect silver to replace hydrogen.

$$M_g(s) + Z$$

$$n SOy(ag) \rightarrow Mg SOy(ag) + Zn(s)$$

magnesium is MORE ACTIVE than zinc, so it should be able to replace zinc in the compound.

CHEMICAL CALCULATIONS - RELATING MASS AND ATOMS



- While chemical equations are written in terms of ATOMS and MOLECULES, that's NOT how we often measure substances in lab!


- measurements are usually MASS (and sometimes VOLUME), NOT number of atoms or molecules! $\bigwedge_{Na_2 CO_3} Solid$ $\bigwedge_{Hcl} Solution$

... so how do we relate atoms and molecules with things we routinely measure in lab - like grams and milliliters?

THE MOLE CONCEPT

- A "mole" of atoms is 6.022×10^{23} why so big? Because atoms are so small!


- Why - in the metric dominated world of science - do we use such a strange number for quantity of atoms?



THE MOLE CONCEPT

- Why define the mole based on an experimentally-measured number?

- The atomic weight of an element (if you put the number in front of the unit GRAMS) is equal to the mass of ONE MOLE of atoms of that element!



Magnesium (Mg): 24.31 g = the mass of ONE MOLE OF MAGNESIUM ATOMS

- So, using the MOLE, we can directly relate a mass and a certain number of atoms!

RELATING MASS AND MOLES

- Use DIMENSIONAL ANALYSIS (a.k.a "drag and drop")

- Need CONVERSION FACTORS - where do they come from?

- We use ATOMIC WEIGHT as a conversion factor.


$$M_{g} : 24.31 | 24.31 g M_{g} = 1 \mod M_{g}$$

"mol" is the abbreviation for
"mole"
Example: How many moles of atoms are there in 250. g of magnesium metal?
24.3 | g M_{g} = mol M_{g}
$$250 \cdot g M_{g} \times \frac{mol M_{g}}{24.3 | g M_{g}} = 10.3 \mod M_{g}$$

Example: You need 1.75 moles of iron. What mass of iron do you need to weigh out on the balance?

SS.85 g Fe = mol Fe
1.75 mol Fe x
$$\frac{SS.85 g Fe}{mol Fe} = 97.7g$$

WHAT ABOUT COMPOUNDS? FORMULA WEIGHT



Formula weight goes by several names:

- For atoms, it's the same thing as ATOMIC WEIGHT
- For molecules, it's called MOLECULAR WEIGHT
- Also called "MOLAR MASS"

Example: How many grams of ammonium carbonate do we need to weigh out to get 3.65 moles of ammonium carbonate?

First, we need to get the formula of the compound; then we can find the formula weight.

$$\frac{NH_{4}^{+}}{(NH_{4})_{2}(\delta_{3})} = \frac{N!2\times14.01}{H!8\times1.008}$$

$$\frac{NH_{4}^{+}}{(NH_{4})_{2}(\delta_{3})} = \frac{N!2\times14.01}{H!8\times1.008}$$

$$\frac{(1\times12.01)}{96.094}$$

$$\frac{0:3\times16.00}{96.094}$$
Formula weight of ammonium carbonate

96.094 g (NH4)2(03 = mo)(NH4)2(03
3.65 mol (NH4)2(03 x
$$\frac{96.094 g (NH4)2(03}{mol(NH4)2(03)} = 351g (NH4)2(03)$$

PERCENTAGE COMPOSITION

- sometimes called "percent composition" or "percent composition by mass"
- the percentage of each element in a compound, expressed in terms of mass Example: Find the percentage composition of ammonium nitrate. NHL NOg-

$$\frac{NH_{4}NO_{3}}{N!2} : N: 2 \times 14.01 = 26.02$$

$$H: 4 \times 1.008 = 4.032$$

$$O: 3 \times 16.00 = \frac{48.00}{80.0529} NH_{4}NO_{3} = mol NH_{4}NO_{3}$$

$$\frac{6}{6}N: \frac{28.029}{80.0529} NH_{4}NO_{3} \times 100\% = 35.0\% N$$

$$\frac{6}{6}H: \frac{4.0529}{80.0529} NH_{4}NO_{3} \times 100\% = 5.0\% N$$

$$\frac{6}{6}O: \frac{48.0090}{80.0529} NH_{4}NO_{3} \times 100\% = 60.0\% N$$

$$\frac{6}{6}O: \frac{48.0090}{80.0529} NH_{4}NO_{3} \times 100\% = 60.0\% 0$$

roundoff

So far, we have



p 483 -

488

- looked at how to determine the composition by mass of a compound

from a formula


- converted from MASS to MOLES (related to the number of atoms/molecules)
- converted from MOLES to MASS

Are we missing anything?

- Sec What about SOLUTIONS, where the desired chemical is not PURE, but found DISSOLVED IN WATER?
 - How do we deal with finding the moles of a desired chemical when it's in solution?

MOLAR CONCENTRATION

- unit: MOLARITY (M): moles of dissolved substance per LITER of solution



There are 6.0 moles of hydrochloric acid in each liter of this solution, so you can write this relationship another way:

6.0 mol HC1 = 1 L

If you have 0.250 L (250 mL) of <u>6.0 M</u> HCI, how many moles of HCI do you have?

6.0 mol HCl =
$$L$$

0.2SOL x $\frac{6.0 \text{ mol HCl}}{L}$ = 1.5 mol HCl

137

If you need 0.657 moles of hydrochloric acid, how many milliliters of 0.0555 M HCl do you need to measure out?

0.0555 mol HC1=L mL=10-3L

O, 657 mol HCl
$$\chi \frac{L}{0.0555}$$
 mol HCl $\chi \frac{mL}{10^{-3}L} = [1800 mL of 0.0555 mHC]$

This is an extremely large volume for lab-scale work (where typical reagent bottles may be 500 mL). We should use a more concentrated solution ... like the 6.00 M HCl below.

What if we used 6.00 M HCl? $6.00 \text{ mol } \text{HCl} = L \text{ m } L = 10^{-3} \text{ L}$

$$O, 657 \text{ mol} HC|_{X} \frac{L}{6.00 \text{ mol} HC|_{X}} \frac{\text{mL}}{10^{-3} \text{L}} = [110 \text{ mL of } 6.00 \text{ mHC}]$$

This volume is much more reasonable than the other - for laboratory work. We can measure this easily with a 250 mL cylinder or we can use a standard 100 mL cylinder twice. If you're preparing a solution by dissolving a solid in water, you can easily calculate the molarity of the solution. How?


Just find the number of moles of solid you dissolved, then divide by the volume of the solution (expressed in liters!)

What is the molarity of a solution made by dissolving 3.50 g of NaCl in enough water to make 250. mL of solution?

$$M = \frac{mol NaCl}{L solution}$$

1 - Find mol NaCl by changing 3.50 g to moles (Use FORMULA WEIGHT)

2 - Divide the mol NaCI and the volume after converting volume to LITERS



A few more examples...

139

Vou have a 250.g bottle of silver(1) chloride (AgCl). How many moles of AgCl do you have? Ag (1 : Ag : [x | 07.9 C1 : 1x 35.45 143.35 g Ag(1 = mol Ag(1)250.g Ag(1 x $\frac{mol Ag(1)}{143.35 g Ag(1)} = \frac{1.74 mol Ag(1)}{1.74 mol Ag(1)}$

How many moles of NaOH are present in 155 mL of 1.50 M NaOH? When relating moles and VOLUME, we need to use CONCENTRATION (usually MOLARITY - M)

$$ISS mL \times \frac{10^{-3}L}{mL} \times \frac{1.50 \text{ mol } NaOH}{L} = 0.233 \text{ mol } NaOH$$

End of material for test 3

Summer 2011 Test 3 is 7/19/2011 (T)