

- thermodynamics: the study of energy transfer

Conservation of energy: Energy may change form, but the overall amount of energy remains constant. "first law of thermodynamics"

- ... but what IS energy?

- energy is the ability to do "work"

↑
motion of matter


Kinds of energy?

- Kinetic energy: energy of matter in motion $E_K = \frac{1}{2} m v^2$

mass

↑
velocity

- Potential energy: energy of matter that is being acted on by a field of force (like gravity)



When the ball falls, its potential energy is converted to kinetic!

- What sort of energy concerns chemists? Energy that is absorbed or released during chemical reactions.

- Energy can be stored in chemicals ... molecules and atoms.

INTERNAL ENERGY: "U"



related to the kinetic and potential energy of atoms, molecules, and their component parts.


- We measure energy transfer ... which is called HEAT. (HEAT is the flow of energy from an area of higher temperature to an area of lower temperature)

Q: heat




SYSTEM: the object or material under study

SURROUNDINGS: everything else


| Type of process | Energy is ... | Sign of Q | Temp of SURROUNDINGS ... |
|-----------------|---|-----------|--------------------------|
| ENDOTHERMIC | transferred from SURROUNDINGS to SYSTEM | + | decreases |
| EXOTHERMIC | transferred from SYSTEM to SURROUNDINGS | - | increases |



The temperature of the flask and the water it contains rises from 25 degrees to 40 degrees. The reaction is EXOTHERMIC, it loses energy to the surroundings (the water in the flask, the flask, the thermometer, etc.) $Q < 0$




Temperature of the flask decreases from 25 degrees to less than 0 degrees. The reaction is ENDOTHERMIC - it absorbs energy from the flask, the hand holding the flask, etc. (the surroundings). $Q > 0$



ENERGY UNITS

- calorie (cal): the amount of energy required to change the temperature of one gram of water by one degree Celsius (or Kelvin)



$1\text{g} \approx 1\text{mL}$ for water

- Calories in food? The "Calorie" that is given on American food labels is actually the kilocalorie (kcal)

- Joule (J): SI unit for energy. It's defined based on the equation for kinetic energy.

$$1\text{ J} = 1 \frac{\text{Kg m}^2}{\text{s}^2}, \text{ from}$$

$$E_K = \frac{1}{2} m v^2$$

kinetic
energy

mass


velocity

$$4.184\text{ J} = 1\text{ cal}$$

- the Joule is a small unit. For most reactions at lab scale, we'll use kilojoules (kJ).

CALORIMETRY

- the measurement of heat. How do we measure heat flow?



... what is Q for this reaction?

Assuming that no heat is lost from the water to the surrounding air,

$$\underbrace{Q_r}_{\text{reaction}} + \underbrace{Q_w}_{\text{water}} = 0$$

Conservation of energy. The terms add to zero because they have opposite signs.

... if we knew something about the WATER, we could use that to find the heat of the REACTION!

SPECIFIC HEAT

- a measured quantity. The amount of energy required to change the temperature of one gram of a particular substance by one degree Celsius.
- Specific heat information for common substances is readily available. For water,

$$4.184 \frac{\text{J}}{\text{g}^\circ\text{C}} \quad \underline{\underline{=}} \quad 1.000 \frac{\text{cal}}{\text{g}^\circ\text{C}}$$

$$Q = m \times s \times \Delta T$$

m = mass
 s = specific heat
 ΔT = $T_{\text{final}} - T_{\text{initial}}$

This is ALWAYS final temp minus initial temp!


- For objects, like reaction vessels, you might know the HEAT CAPACITY, which is the amount of energy required to change the temperature of an object by one degree Celsius

$$\text{units: } \text{J}/^\circ\text{C} \quad \text{or} \quad \text{cal}/^\circ\text{C}$$


$$Q = C \times \Delta T$$

c = heat capacity

0.20 mol A



When we add the reactant to water, it decomposes - heating the water.



Specific heat of water:

$$4.184 \frac{\text{J}}{\text{g}^\circ\text{C}}$$

$$Q_r + Q_w = 0$$

$$\begin{aligned} Q_w &= m_w \times S_w \times \Delta T_w \\ &= (100 \text{ g}) (4.184 \frac{\text{J}}{\text{g}^\circ\text{C}}) (38^\circ\text{C} - 25^\circ\text{C}) \\ &= 5439.2 \text{ J} \end{aligned}$$

$$Q_r + 5439.2 \text{ J} = 0; Q_r = -5439.2 \text{ J}$$

To report the energy change in this reaction to others, we should express it in terms of heat transfer per mole of something. A different amount of reactant would have a different Q

$$Q_{\text{rxn}} = \frac{Q_r}{\text{moles A}} = \frac{-5439.2 \text{ J}}{0.20 \text{ mol A}} = -27000 \frac{\text{J}}{\text{mol A}} = \boxed{-27 \frac{\text{kJ}}{\text{mol A}}}$$

This number is often called the "HEAT OF REACTION"!

One problem ...

PATH. The amount of energy required for a process depends on how the process is carried out.

Example: Driving from Florence to Columbia. How much energy is required? (gas)

2000 Jeep Cherokee vs 2008 Toyota Prius. The Jeep will use much more fuel than the Prius even though they start and end from exactly the same place. So the fuel usage is what we call a PATH FUNCTION, while the location is a STATE FUNCTION.

- so the heat of reaction depends on how the reaction is done.
- we need (for reporting) some kind of standard condition. At constant pressure, we can define a state function called ENTHALPY (H)

$$H = U + PV$$

$$\Delta H = Q_{\text{constant pressure}}$$

... we record the "enthalpy change of reaction" in our data books.

$$\Delta H_r$$