


<sup>158</sup> SINCE the enthalpy change does NOT depend on path, this means that we can use standard values for enthalpy to predict the heat change in reactions that we have not tested in a calorimeter.

---

## THERMOCHEMICAL EQUATIONS

- is like a regular chemical equation, except that phase labels are REQUIRED and the enthalpy for the reaction is given along with the equation.



- Why are phase labels required? Because phase changes either absorb or release energy.

$\Delta H = -1800 \text{ kJ}$  ... what does this mean?

$$1 \text{ mol CH}_3\text{COCH}_3 = -1800 \text{ kJ}$$

$$4 \text{ mol O}_2 = -1800 \text{ kJ}$$

$$3 \text{ mol CO}_2 = -1800 \text{ kJ}$$

$$3 \text{ mol H}_2\text{O} = -1800 \text{ kJ}$$

We treat the enthalpy change as if it's another product of the reaction!



What would be the enthalpy change when 25 g of water are produced by the reaction?

- 1 - Convert 25.0 g water to moles using formula weight.
- 2 - Convert moles water to enthalpy change using thermochemical equation.

$$18.016 \text{ g H}_2\text{O} = 1 \text{ mol H}_2\text{O} \quad | \quad 3 \text{ mol H}_2\text{O} = -1800 \text{ kJ}$$

$$25.0 \text{ g H}_2\text{O} \times \frac{1 \text{ mol H}_2\text{O}}{18.016 \text{ g H}_2\text{O}} \times \frac{-1800 \text{ kJ}}{3 \text{ mol H}_2\text{O}} = \boxed{-830 \text{ kJ}}$$

- This reaction is EXOTHERMIC
- Energy is released from the reaction mixture, heating the surroundings. (This is true for all combustions...)
- Provided that the reaction takes place at constant pressure, this enthalpy change should be equal to Q (measurable).

A few more terms related to enthalpy:


- Enthalpy of vaporization / heat of vaporization: The enthalpy change on vaporizing one mole of a substance. (from liquid to vapor)
- Enthalpy of fusion / heat of fusion: The enthalpy change when a mole of liquid changes to the solid state.




Phase changes require energy, too!

## 160 FORMATION REACTIONS

- A reaction that forms exactly one mole of the specified substance from its elements at their STANDARD STATE at 25C and 1 atm pressure.



heat of formation of carbon dioxide  $\Delta H_f^\circ$  or  $\Delta H_f$   
"enthalpy of formation"



you may see fractional coefficients in these formation reactions, because you MUST form exactly one mole of the product!


- The heat of formation for an element in its standard state at 25C and 1 atm is ZERO.

$$\Delta H_f^\circ, \text{O}_2(g) = 0 \text{ kJ/mol}$$

- What are formation reactions good for?





... finding enthalpies for more interesting reactions!

<sup>161</sup> Let's say we would like to find the enthalpy of reaction for this equation:



From  
A-8,  
text  
↓

Hess' Law: If you add two reactions to get a new reaction, their enthalpies also add.



$$\Delta H = 2(-226.7) + 4(-393.5) + 2(-241.8) = \boxed{-2511 \text{ kJ}}$$


$$\Delta H = \sum \Delta H_{f, \text{products}} - \sum \Delta H_{f, \text{reactants}}$$

$$\Delta H = 4(-393,5) + 2(-241,8) - (2(226,7) + 5(0))$$
$$= 2511 \text{ kJ}$$

See Appendix C in the textbook for enthalpy of formation data:  
p A-8 to A-11

\* Remember:

- Multiply each enthalpy by its stoichiometric coefficient from the reaction
- Enthalpy of formation of an element at its standard state is zero
- Watch phase labels. You will usually find SEVERAL enthalpies of formation for a given substance in different phases!
- For ionic substances in solution, remember that they exist as free ions, so look up the aqueous IONS!



Calculate the enthalpy change for the combustion of 1.00 kg of hydrogen gas.

1 - Convert 1.00 kg of hydrogen gas to moles using formula weight.

2 - Convert moles hydrogen gas to enthalpy change using thermochemical equation.

$$2.016 \text{ g H}_2 = 1 \text{ mol H}_2 \quad | \quad 2 \text{ mol H}_2 = -484 \text{ kJ} \quad | \quad \text{kg} = 10^3 \text{ g}$$

$$1.00 \text{ kg H}_2 \times \frac{10^3 \text{ g}}{\text{kg}} \times \frac{1 \text{ mol H}_2}{2.016 \text{ g H}_2} \times \frac{-484 \text{ kJ}}{2 \text{ mol H}_2} = \boxed{-120000 \text{ kJ per kg H}_2}$$