

If 50.3 kJ of heat was released when 5.48 g of formic acid are burned at constant pressure, then what is the enthalpy change of this reaction per mole of formic acid?


$$Q = -50.3 \text{ kJ}; \Delta H = \frac{Q_{\text{constant pressure}}}{\text{mol HCHO}_2}$$

Find moles formic acid:


$$5.48 \text{ g HCHO}_2 \times \frac{\text{mol HCHO}_2}{46.026 \text{ g HCHO}_2} = 0.1190631382 \text{ mol HCHO}_2$$

$$\Delta H = \frac{Q_{\text{constant pressure}}}{\text{mol HCHO}_2} = \frac{-50.3 \text{ kJ}}{0.1190631382 \text{ mol HCHO}_2} = \boxed{-422 \text{ kJ/mol HCHO}_2}$$

Based on the calculation above, can we complete this thermochemical equation?



We calculated the enthalpy change per mole of formic acid in the calorimetry problem above, and this equation is based on TWO MOLES of formic acid. So we double the enthalpy change per mole to get the change for this reaction.



What is the enthalpy change when 150. L of nitrogen monoxide are formed by this reaction at 25.0 C and 1.50 atm pressure?

1 - Convert 150. L NO to moles using ideal gas equation.


2 - Convert moles NO to enthalpy change using thermochemical equation.

$$\textcircled{1} \quad PV = nRT \quad \left| \quad \begin{array}{l} P = 1.50 \text{ atm} \quad R = 0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}} \\ n = \frac{PV}{RT} \quad \left| \quad \begin{array}{l} V = 150. \text{ L} \\ T = 25.0^\circ\text{C} = 298.2 \text{ K} \end{array} \right. \end{array} \right.$$

$$n_{\text{NO}} = \frac{(1.50 \text{ atm})(150. \text{ L})}{\left(0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}}\right)(298.2 \text{ K})} = 9.194822849 \text{ mol NO}$$

$$\textcircled{2} \quad 4 \text{ mol NO} = -906 \text{ kJ}$$

$$9.194822849 \text{ mol NO} \times \frac{-906 \text{ kJ}}{4 \text{ mol NO}} = \boxed{-2080 \text{ kJ}}$$



What is the enthalpy change at standard conditions when 25.0 grams of hydrogen sulfide gas is reacted?

- 1 - Calculate the enthalpy change for the reaction as written using heats of formation.
- 2 - Convert 25.0 g hydrogen sulfide to moles using formula weight
- 3 - Convert moles hydrogen sulfide to enthalpy change using thermochemical equation

$$\begin{aligned} \textcircled{1} \Delta H &= \sum \Delta H_f^\circ, \text{ products} - \sum \Delta H_f^\circ, \text{ reactants} \\ &= [2(-285.8) + 2(-296.8)] - [2(-20.50) + 3(0)] = -1124.2 \text{ kJ} \end{aligned}$$

Thermochemical equation:




34.086 g H ₂ S = mol H ₂ S	2 mol H ₂ S = -1124.8 kJ
--------------------------------------------------	-------------------------------------

$$25.0 \text{ g H}_2\text{S} \times \frac{\text{mol H}_2\text{S}}{34.086 \text{ g H}_2\text{S}} \times \frac{-1124.8 \text{ kJ}}{2 \text{ mol H}_2\text{S}} = \boxed{-412 \text{ kJ}}$$

②
②

propane



Calculate the volume of propane gas at 25.0 C and 1.08 atm required to provide 565 kJ of heat using the reaction above.

- 1 - Convert energy requirement to moles of PROPANE using thermochemical equation
- 2 - Convert moles propane to volume using ideal gas equation

① $\text{mol C}_3\text{H}_8 = -2043 \text{ kJ}$ Since the reaction is providing the heat, the heat is given a negative sign. From the point of view of the reaction, it's an EXOTHERMIC process!

$$-565 \text{ kJ} \times \frac{\text{mol C}_3\text{H}_8}{-2043 \text{ kJ}} = 0.2765540671 \text{ mol C}_3\text{H}_8$$



$$\begin{array}{l|l} PV = nRT & n = 0.2765540671 \text{ mol C}_3\text{H}_8 \quad T = 25.0^\circ\text{C} = 298.2 \text{ K} \\ V = \frac{nRT}{P} & R = 0.08206 \frac{\text{L}\cdot\text{atm}}{\text{mol}\cdot\text{K}} \quad P = 1.08 \text{ atm} \end{array}$$

$$V = \frac{(0.2765540671 \text{ mol C}_3\text{H}_8) \left(0.08206 \frac{\text{L}\cdot\text{atm}}{\text{mol}\cdot\text{K}}\right) (298.2 \text{ K})}{(1.08 \text{ atm})}$$

$$= 6.27 \text{ L propane at } 25.0^\circ\text{C}, 1.08 \text{ atm}$$

END OF CHAPTER 6

LIGHT



Diffraction

frequency = wavelengths / time = ν s^{-1} : Hertz, Hz

- Light has properties of WAVES such as DIFFRACTION (it bends around small obstructions).
- Einstein noted that viewing light as a particle that carried an energy proportional to the FREQUENCY could explain the PHOTOELECTRIC EFFECT!


$$E_{\text{photon}} = h \nu$$

Planck's constant: $6.63 \times 10^{-34} \text{ J}\cdot\text{s}$

photon = particle or packet of light

(The photoelectric effect is the emission of electrons from a metal caused by exposure to light. Einstein discovered that if the light were not of the correct FREQUENCY, increasing the INTENSITY of the light would not cause electron emission. He concluded that individual photons must have enough energy to excite an electron - i.e. they must have the appropriate frequency.)

The photoelectric effect and Einstein's ideas about the energy content of light led us to discover a new model for the atom! How? Let's start with the nuclear model:



Nuclear model:

- Protons and neutrons in a dense NUCLEUS at center of atom
- Electrons in a diffuse (mostly empty) ELECTRON CLOUD surrounding NUCLEUS.

... so what's wrong with the nuclear model? Among other things, it doesn't explain ...