

## PERCENT YIELD

- Chemical reactions do not always go to completion! Things may happen that prevent the conversion of reactants to the desired/expected product!


① SIDE REACTIONS:



... so in a low-oxygen environment, you may produce less carbon dioxide than expected!

② TRANSFER AND OTHER LOSSES

- When isolating a product, losses may occur in the process. Example: filtering



### ③ EQUILIBRIUM

- Reactions may reach an equilibrium between products and reactants. We'll talk more about this in CHM 111. The net result is that the reaction will appear to stop before all reactants have been consumed!


- All of these factors cause a chemical reaction to produce LESS product than calculated. For many reactions, this difference isn't significant. But for others, we need to report the PERCENT YIELD.

$$\text{PERCENT YIELD} = \frac{\text{ACTUAL YIELD}}{\text{THEORETICAL YIELD}} \times 100\%$$

↙ Determined EXPERIMENTALLY!

↑ Calculated based on the limiting reactant. (The chemical calculations you've done up to now have been theoretical yields!)

... the percent yield of a reaction can never be greater than 100% due to conservation of mass! If you determine that a percent yield is greater than 100%, then you've made a mistake somewhere - either in a calculation or in the experiment itself!



22.4 grams of benzene are reacted with excess nitric acid. If 31.6 grams of nitrobenzene are collected from the reaction, what is the percent yield?

To determine the PERCENT YIELD, we need to calculate the THEORETICAL YIELD - the amount of nitrobenzene that COULD be produced from 22.4 grams of benzene.




$$22.4 \text{ g C}_6\text{H}_6 \times \frac{\text{mol C}_6\text{H}_6}{78.114 \text{ g C}_6\text{H}_6} \times \frac{\text{mol C}_6\text{H}_5\text{NO}_2}{\text{mol C}_6\text{H}_6} \times \frac{123.111 \text{ g C}_6\text{H}_5\text{NO}_2}{\text{mol C}_6\text{H}_5\text{NO}_2} = 35.3 \text{ g C}_6\text{H}_5\text{NO}_2$$

THEORETICAL YIELD

$$\text{PERCENT YIELD} = \frac{\text{ACTUAL YIELD}}{\text{THEOR. YIELD}} \times 100\% = \frac{31.6 \text{ g}}{35.3 \text{ g}} \times 100\% = \boxed{89.5\%}$$

25.0 mL of acetic acid solution requires 37.3 mL of 0.150 M sodium hydroxide for complete reaction. The equation for this reaction is:



What is the molar concentration of the acetic acid?

$$\frac{\text{L mol HC}_2\text{H}_3\text{O}_2}{\text{L solution}} \leftarrow = 25.0\text{mL or } 0.0250\text{L}$$


Since we already know the volume of the acetic acid solution, we just need to calculate the MOLES OF ACETIC ACID in the solution. Then, we can find molarity by division.

$$m\text{L} = 10^{-3}\text{L} \quad 0.150\text{ mol NaOH} = \text{L} \quad \text{mol NaOH} = \text{mol HC}_2\text{H}_3\text{O}_2$$

$$37.3\text{mL} \times \frac{10^{-3}\text{L}}{\text{mL}} \times \frac{0.150\text{ mol NaOH}}{\text{L}} \times \frac{\text{mol HC}_2\text{H}_3\text{O}_2}{\text{mol NaOH}} = 0.005595\text{ mol HC}_2\text{H}_3\text{O}_2$$

Now, find molarity:

$$M = \frac{\text{mol HC}_2\text{H}_3\text{O}_2}{\text{L solution}} = \frac{0.005595\text{ mol HC}_2\text{H}_3\text{O}_2}{0.0250\text{L}} = \boxed{0.224\text{ M HC}_2\text{H}_3\text{O}_2}$$



Calculate how many grams of acrylonitrile could be obtained from 651 kg of propylene, assuming there is excess NO present.

- 1 - Convert 651 kg of propylene to moles propylene using formula weight. (kg -> g conversion)
- 2 - Convert moles propylene to moles acrylonitrile using chemical equation.
- 3 - Convert moles acrylonitrile to mass using formula weight.

$$42.081 \text{ g C}_3\text{H}_6 = 1 \text{ mol C}_3\text{H}_6 \quad | \quad 4 \text{ mol C}_3\text{H}_6 = 4 \text{ mol C}_3\text{H}_3\text{N} \quad | \quad \text{kg} = 10^3 \text{ g}$$


$$53.064 \text{ g C}_3\text{H}_3\text{N} = 1 \text{ mol C}_3\text{H}_3\text{N}$$

$$651 \text{ kg C}_3\text{H}_6 \times \frac{10^3 \text{ g}}{\text{kg}} \times \frac{1 \text{ mol C}_3\text{H}_6}{42.081 \text{ g C}_3\text{H}_6} \times \frac{4 \text{ mol C}_3\text{H}_3\text{N}}{4 \text{ mol C}_3\text{H}_6} \times \frac{53.064 \text{ g C}_3\text{H}_3\text{N}}{1 \text{ mol C}_3\text{H}_3\text{N}} =$$

①
②
③

$$= 821000 \text{ g C}_3\text{H}_3\text{N} \quad (821 \text{ kg})$$

151.90 g/mol



How many mL of 0.250M potassium permanganate are needed to react with 3.36 g of iron(II) sulfate?

- 1 - Convert mass iron(II) sulfate to moles using formula weight.
- 2 - Convert moles iron(II) sulfate to moles potassium permanganate using chemical equation
- 3 - Convert moles potassium permanganate to volume using concentration

$$151.90 \text{ g FeSO}_4 = \text{mol FeSO}_4 \quad | \quad 10 \text{ mol FeSO}_4 = 2 \text{ mol KMnO}_4$$

$$0.250 \text{ mol KMnO}_4 = \text{L} \quad | \quad \text{mL} = 10^{-3} \text{ L}$$

$$3.36 \text{ g FeSO}_4 \times \frac{\text{mol FeSO}_4}{151.90 \text{ g FeSO}_4} \times \frac{2 \text{ mol KMnO}_4}{10 \text{ mol FeSO}_4} \times \frac{\text{L}}{0.250 \text{ mol KMnO}_4} \times \frac{\text{mL}}{10^{-3} \text{ L}} =$$

①
②
③

$$= 17.7 \text{ mL of } 0.250 \text{ M KMnO}_4$$

## Electrolytes and Ionic Theory

- electrolytes: substances that dissolve in water to form charge-carrying solutions

\* Electrolytes form ions in solution - (ions that are mobile are able to carry charge!). These IONS can undergo certain kinds of chemistry!

---

### IONIC THEORY

- the idea that certain compounds DISSOCIATE in water to form free IONS

What kind of compounds?

- Soluble ionic compounds
- Acids (strong AND weak)
- Bases (strong AND weak)

The ions formed may interact with each other to form NEW compounds!

Strong vs weak?

- If an electrolyte COMPLETELY IONIZES in water, it's said to be STRONG
- If an electrolyte only PARTIALLY IONIZES in water, it's said to be WEAK
- Both kinds of electrolyte undergo similar kinds of chemistry.