


How many milliliters of 6.00M hydrochloric acid is needed to completely react with 25.0 g of sodium carbonate?



- 1 - Convert 25.0 g of sodium carbonate to moles. Use FORMULA WEIGHT.
- 2 - Convert moles sodium carbonate to moles HCl. Use CHEMICAL EQUATION.
- 3 - Convert moles HCl to volume. Use MOLAR CONCENTRATION of HCl (and a L→mL conversion)


$$\textcircled{3} \quad 6.00 \text{ mol HCl} = \text{L} \quad | \quad \text{mL} = 10^{-3} \text{ L}$$


$$0.4717426172 \text{ mol HCl} \times \frac{\text{L}}{6.00 \text{ mol HCl}} \times \frac{\text{mL}}{10^{-3} \text{ L}} = 78.6 \text{ mL of } 6.00\text{M HCl}$$


The problem asked for the volume in milliliter units, so we needed to convert from L → mL after using molarity.

CONCEPT OF LIMITING REACTANT

- When does a chemical reaction STOP?




Magnesium
strip


Flame,
oxygen from
air


Magnesium oxide
powder

- When does this reaction stop? When burned in open air, this reaction stops when all the MAGNESIUM STRIP is gone. We say that the magnesium is LIMITING.

- This reaction is controlled by the amount of available magnesium


- At the end of a chemical reaction, the LIMITING REACTANT will be completely consumed, but there may be amount of OTHER reactants remaining. We do chemical calculations in part to minimize these "leftovers".

 These are often called "excess" reactants, or reactants present "in excess"

LIMITING REACTANT CALCULATIONS

- To find the limiting reactant, calculate how much product would be produced from ALL given reactants. Whichever produces the SMALLEST amount of product is the limiting reactant, and the smallest amount of product is the actual amount of product produced.

Example: 56.08 12.01 64.10 <- Formula weights



If you start with 100. g of each reactant, how much calcium carbide would be produced?

$$\text{CaO: } 56.08 \text{ g CaO} = \text{mol CaO} \quad | \quad \text{mol CaO} = \text{mol CaC}_2 \quad | \quad 64.10 \text{ g CaC}_2 = \text{mol CaC}_2$$

$$100. \text{ g CaO} \times \frac{\text{mol CaO}}{56.08 \text{ g CaO}} \times \frac{\text{mol CaC}_2}{\text{mol CaO}} \times \frac{64.10 \text{ g CaC}_2}{\text{mol CaC}_2} = 114 \text{ g CaC}_2$$

$$\text{C: } 12.01 \text{ g C} = \text{mol C} \quad | \quad 3 \text{ mol C} = \text{mol CaC}_2 \quad | \quad 64.10 \text{ g CaC}_2 = \text{mol CaC}_2$$

$$100. \text{ g C} \times \frac{\text{mol C}}{12.01 \text{ g C}} \times \frac{\text{mol CaC}_2}{3 \text{ mol C}} \times \frac{64.10 \text{ g CaC}_2}{\text{mol CaC}_2} = 178 \text{ g CaC}_2$$

114 grams of calcium carbide should be produced.. When 114 grams of the carbide is formed, all 100 grams of CaO have been used up, and the reaction stops. We still have leftover carbon, but there's nothing for it to react with! - no more product can be made!

We say that CaO is "limiting" (it controls how much product we make), and C is present "in excess".

PERCENT YIELD

- Chemical reactions do not always go to completion! Things may happen that prevent the conversion of reactants to the desired/expected product!


① SIDE REACTIONS:



... so in a low-oxygen environment, you may produce less carbon dioxide than expected!

② TRANSFER AND OTHER LOSSES

- When isolating a product, losses may occur in the process. Example: filtering



③ EQUILIBRIUM

- Reactions may reach an equilibrium between products and reactants. We'll talk more about this in CHM 111. The net result is that the reaction will appear to stop before all reactants have been consumed!


- All of these factors cause a chemical reaction to produce LESS product than calculated. For many reactions, this difference isn't significant. But for others, we need to report the PERCENT YIELD.

$$\text{PERCENT YIELD} = \frac{\text{ACTUAL YIELD}}{\text{THEORETICAL YIELD}} \times 100\%$$

↙ Determined EXPERIMENTALLY!

↑ Calculated based on the limiting reactant. (The chemical calculations you've done up to now have been theoretical yields!)

... the percent yield of a reaction can never be greater than 100% due to conservation of mass! If you determine that a percent yield is greater than 100%, then you've made a mistake somewhere - either in a calculation or in the experiment itself!



22.4 grams of benzene are reacted with excess nitric acid. If 31.6 grams of nitrobenzene are collected from the reaction, what is the percent yield?

To get percent yield, we need to calculate the THEORETICAL YIELD of nitrobenzene, starting from the 22.4 g of benzene we reacted.


$$78.114 \text{ g C}_6\text{H}_6 = \text{mol C}_6\text{H}_6 \quad | \quad \text{mol C}_6\text{H}_6 = \text{mol C}_6\text{H}_5\text{NO}_2$$

$$123.111 \text{ g C}_6\text{H}_5\text{NO}_2 = \text{mol C}_6\text{H}_5\text{NO}_2$$

$$\begin{aligned}
 22.4 \text{ g C}_6\text{H}_6 & \times \frac{\text{mol C}_6\text{H}_6}{78.114 \text{ g C}_6\text{H}_6} \times \frac{\text{mol C}_6\text{H}_5\text{NO}_2}{\text{mol C}_6\text{H}_6} \times \frac{123.111 \text{ g C}_6\text{H}_5\text{NO}_2}{\text{mol C}_6\text{H}_5\text{NO}_2} = \\
 & = 35.3 \text{ g C}_6\text{H}_5\text{NO}_2 \text{ (theoretical yield)}
 \end{aligned}$$

$$\text{Percent yield} = \frac{\text{actual yield}}{\text{theor. yield}} \times 100\% = \frac{31.6 \text{ g}}{35.3 \text{ g}} \times 100\% = \boxed{89.5\%}$$

25.0 mL of acetic acid solution requires 37.3 mL of 0.150 M sodium hydroxide for complete reaction. The equation for this reaction is:



What is the molar concentration of the acetic acid?

$$\frac{\text{L mol HC}_2\text{H}_3\text{O}_2}{\text{L solution} \leftarrow = 25.0\text{mL or } 0.0250\text{L}}$$

Since we already know the volume of acetic acid solution, we need to find the moles of acetic acid. To do THAT, we'll start with the sodium hydroxide volume, since we can relate that to moles (and then to the amount of acetic acid)!

$$\text{mL} = 10^{-3} \text{ L} \quad 0.150 \text{ mol NaOH} = \text{L} \quad \text{mol NaOH} = \text{mol HC}_2\text{H}_3\text{O}_2$$

$$37.3 \text{ mL} \times \frac{10^{-3} \text{ L}}{\text{mL}} \times \frac{0.150 \text{ mol NaOH}}{\text{L}} \times \frac{\text{mol HC}_2\text{H}_3\text{O}_2}{\text{mol NaOH}} = 0.005595 \text{ mol HC}_2\text{H}_3\text{O}_2$$

To get CONCENTRATION, we divide the moles by the volume of the acetic acid (in L)

$$M = \frac{\text{mol HC}_2\text{H}_3\text{O}_2}{\text{L solution}} = \frac{0.005595 \text{ mol HC}_2\text{H}_3\text{O}_2}{0.0250 \text{ L}} = 0.224 \text{ M HC}_2\text{H}_3\text{O}_2$$