Example: How would we prepare 500. mL of 0.500 M sodium sulfate in water?

Naz S04: 142.05 g/mol

H20

Dissolve the appropriate amount of sodium sulfate into enough water to make 500. mL of

solution.



A VOLUMETRIC FLASK is a flask that is designed to precisely contain a certain volume of liquid.

VOLUMETRIC FLASKS are used to prepare solutions.

volumetric flask

We know that we need 500. mL of solution, and we also know that the concentration should be 0.500 M. From that, we can calculate the moles of sodium sulfate we should dissolve. Then, we can convert that to mass using formula weight.

So, to prepare this solution, we will add 35.5 grams of sodium sulfate to a 500. mL volumetric flask, and dilute to the mark with distilled or deionized water.

More on MOLARITY

To prepare a solution of a given molarity, you generally have two options:

- Weigh out the appropriate amount of solute, then dilute to the desired volume with solvent (usually water)
- Take a previously prepared solution of known concentration and DILUTE it with solvent to form a new solution

- Use DILUTION EQUATION

The dilution equation is easy to derive with simple algebra.

... but when you dilute a solution, the number of moles of solute REMAINS CONSTANT. (After all, you're adding only SOLVENT)

$$M_1 V_1 = M_2 V_2$$
 Since the number of moles of solute stays before after the same, this equality must be true!

$$M_1 V_1 = M_2 V_2$$
 ... the "DILUTION EQUATION" $M_1 = \text{molarity of concentrated solution}$

 $\sqrt{1 - volume}$ of concentrated solution

 M_2 molarity of dilute solution

The volumes don't HAVE to be in liters, as long as you use the same volume UNIT for both volumes!

Example: Take the 0.500 M sodium sulfate we discussed in the previous example and dilute it to make 150. mL of 0.333 M solution. How many mL of the original solution will we need to dilute?

$$M_1 V_1 = M_2 V_2$$

 $(0.500 \, \text{m}) V_1 = (0.333 \, \text{m}) (150. \, \text{mL})$
 $V_1 = 99.9 \, \text{mL of 0.500 M Now, Soy}$

So, we would take 99.9 mL of 0.500 M sodium sulfate, and add enough water to make 150. mL of solution. (Ideally, do this in a 150 mL volumetric flask, if you have one)

- Chemical reactions proceed on an ATOMIC basis, NOT a mass basis!
- To calculate with chemical reactions (i.e. use chemical equations), we need everything in terms of ATOMS ... which means MOLES of atoms

2 All(s) +3 Br₂(l)
$$\rightarrow$$
 2 Al Br₃(s)

Toefficients are in terms of atoms and molecules!

2 atoms Al = 3 molecules Br₂ = 2 formula units Al Br₃

2 mol Al = 3 mol Br₂ = 2 mol Al Br₃

- To do chemical calculations, we need to:
 - Relate the amount of substance we know (mass or volume) to a number of moles
 - Relate the moles of one substance to the moles of another using the equation
 - Convert the moles of the new substance to mass or volume as desired

$$2A(ls) + 3Br_2(l) \longrightarrow 2A(Br_3(s))$$

- * Given that we have 25.0 g of liquid bromine, how many grams of aluminum would we need to react away all of the bromine? How many grams of aluminum bromide would be produced?
 - Convert grams of bromine to moles: Need formula weight B_{12} : $\frac{2 \times 79.90}{159.80}$ $\frac{159.80}{25.09BC_2} \times \frac{1 \text{ mol } BC_2}{159.80gBC_2} = 0.15645 \text{ mol } BC_2$
 - Use the chemical equation to relate moles of bromine to moles of aluminum 2 mol A = 3 mol BG

3 Convert moles aluminum to mass: Need formula weight A1:26,78 26,989 A1=1 mol A1

You can combine all three steps on one line if you like!

You can solve the second part of the question using CONSERVATION OF MASS - since there's only a single product and you already know the mass of all reactants.

But ...

...what would you have done to calculate the mass of aluminum bromide IF you had NOT been asked to calculate the mass of

$$25.0 g Br_2 \times \frac{1 mol Br_2}{159.40 g Br_2} \times \frac{2 mol AlBr_3}{3 mol Br_2} \times \frac{266.694 g AlBr_3}{1 mol AlBr_3} = 27.8 g$$

$$\frac{2}{1 mol AlBr_3} \times \frac{266.694 g AlBr_3}{1 mol AlBr_3} = 27.8 g$$

$$\frac{2}{1 mol AlBr_3} \times \frac{2}{1 mol AlBr_3} \times \frac{27.8 g}{1 mol AlBr_3} \times \frac{2}{1 mol AlBr_3} = 27.8 g$$

$$\frac{2}{1 mol AlBr_3} \times \frac{2}{1 mol AlBr_3} \times \frac{2}{1 mol AlBr_3} \times \frac{2}{1 mol AlBr_3} = 27.8 g$$

$$\frac{2}{1 mol AlBr_3} \times \frac{2}{1 m$$

101 Example:

How many milliliters of 6.00M hydrochloric acid is needed to completely react with 25.0 g of sodium carbonate?

$$=2H(1(aq)+Na_2(0_3(s)) \rightarrow H_20(l)+(0_2(g)+2N_1c)(aq)$$

- 1 Convert 25.0 g sodium carbonate to moles. Use FORMULA WEIGHT.
- 2 Convert moles sodium carbonate to moles HCI. Use CHEMICAL EQUATION.
- 3 Convert moles HCI to volume. Use MOLAR CONCENTRATION of HCI (and convert L->mL)

$$0.2358713086 \text{ mol } Na_2(0_3 \times \frac{2 \text{ mol } H(1)}{\text{mol } Na_2(0_3)} = 0.4717426172 \text{ mol } H(1)$$

102 Example:

How many milliliters of 6.00M hydrochloric acid is needed to completely react with 25.0 g of sodium carbonate?

- 1 Convert 25.0 g sodium carbonate to moles. Use FORMULA WEIGHT.
- 2 Convert moles sodium carbonate to moles HCI. Use CHEMICAL EQUATION.
- 3 Convert moles HCI to volume. Use MOLAR CONCENTRATION of HCI (and convert L->mL)

We did this last conversion because the problem specifically asked for an answer in MILLILITERS!