


... but how do we use this to describe a reaction that produces ions? Let's look at our previous example!




... this is a bit easier to follow than looking at all those letters and numbers in the electron configurations for these elements!

MOLECULAR COMPOUNDS


- Form when atoms SHARE electrons instead of transferring them. This results in the formation of MOLECULES ... groups of atoms held together by electron-sharing.

How might atoms SHARE electrons? By coming together close enough so that their atomic ORBITALS overlap each other:

This idea is called
VALENCE
BOND
THEORY!



Each hydrogen atom has a single electron in a 1s orbital.




When hydrogen atoms come close enough to each other for these orbitals to OVERLAP, each hydrogen "sees" BOTH electrons, filling up the "s" orbitals of both atoms. This is a COVALENT BOND.

Co: together
Valent: valence shells

This is the DIATOMIC MOLECULE, H_2

... so how would this look using dot notation?




In dot structures, SHARED PAIRS of electrons are often written as DASHES to make the structures look neater.



* Why doesn't hydrogen end up with eight electrons? Because hydrogen has only the first shell, which contains only a single "s" subshell (NO "p" subshell). This "s" subshell is full with two electrons, and that's all hydrogen needs to get.

Let's look at OXYGEN ...



We know that oxygen exists in air as the diatomic molecule O_2



The oxygen atoms share TWO pairs of electrons. This is called a DOUBLE BOND

OR




Each oxygen atom has a share in eight electrons!


A few notes on the double bond:

- ① - For atoms to share more than one pair of electrons, they have to move closer to one another than they would if they were only sharing one pair of electrons. This BOND DISTANCE is measurable!
- ② - It takes more energy to break a double bond between two atoms than it would to break a single bond between the same two atoms. This BOND ENERGY is also measurable!

Let's look at NITROGEN ...



We know that nitrogen exists in air as the diatomic molecule N_2



The nitrogen atoms share THREE pairs of electrons. This is called a TRIPLE BOND

OR



Nitrogen gas is fairly inert ... it's hard to break the triple bond in nitrogen gas apart!

A few notes on the triple bond:

- ① - For atoms to share three pairs of electrons, they have to move closer to one another than they would if they were sharing one or two pairs of electrons. Triple bonds have the shortest BOND DISTANCE of all covalent bonds.
- ② - It takes more energy to break a triple bond between two atoms than it would to break either a single or double bond between the same two atoms. The triple bond has the largest BOND ENERGY of all three kinds of covalent bonds.

SO FAR, we've seen that ...





- ① Atoms may share one, two, or three pairs of electrons with each other.
- ② Atoms will usually share enough electrons so that each atom ends up with a share in EIGHT electrons - the "octet rule"

- HYDROGEN will only end up with two electrons!

- Some other atoms may end up with more or less than eight electrons. Exceptions to the octet rule are covered in Chapter 9.

NOW, how could we come up with dot structures for some more complicated (and therefore, more interesting) molecules?

Examples:



① Count valence electrons


② Pick central atom and draw skeletal structure

- central atom is usually the one that needs to gain the most electrons!


- skeletal structure has all atoms connected to center with single bonds

③ Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.

④ Check octet rule - each atom should have a share in 8 electrons (H gets 2). if not, make double or triple bonds.



Choose CARBON as central atom, since it needs the most electrons.




Distribute the remaining electrons. Stop here because we've used all 24.



... but this CARBON atom has a share in only SIX electrons!

We need a double bond, but which atom is most likely to double bond? We choose OXYGEN to form the double bond because it needed an additional TWO electrons at the start, and is likely able to share two pairs of electrons - one for each electron needed.



This structure looks better, Each atom has a share in eight electrons!

- Count valence electrons
- Pick central atom and draw skeletal structure

- central atom is usually the one that needs to gain the most electrons!
- skeletal structure has all atoms connected to center with single bonds

- Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.

- Check octet rule - each atom should have a share in 8 electrons (H gets 2). If not, make double or triple bonds.



$$\text{N: } 1 \times 5$$


$$\text{O: } 1 \times 6$$

$$\text{Cl: } 1 \times 7$$

$$18e^-$$



We pick NITROGEN as the central atom, since it needs more electrons than O or Cl.



We ran out of space on the outer atoms, so the last two electrons go onto the central nitrogen.



Using a pair of electrons from oxygen (to make a double bond) "fixes" this structure!

① Count valence electrons



② Pick central atom and draw skeletal structure

- central atom is usually the one that needs to gain the most electrons!

- skeletal structure has all atoms connected to center with single bonds

③ Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.

④ Check octet rule - each atom should have a share in 8 electrons (H gets 2). If not, make double or triple bonds.



... but each carbon has a share in only FOUR electrons.



... now the carbon has a share in SIX electrons.




Adding a second double bond with the other oxygen atom fixes this structure and gives carbon a share in eight electrons.



The two oxygen atoms are in identical environments and SHOULD bond in the same way, not in the arrangement shown to the left.


EXPERIMENTALLY, we can find (via measuring bond distance) that the two oxygen atoms are the same distance away from the central carbon atom. This result doesn't agree with the triple-bond structure above!

- ① Count valence electrons
- ② Pick central atom and draw skeletal structure
 - central atom is usually the one that needs to gain the most electrons!
 - skeletal structure has all atoms connected to center with single bonds
- ③ Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.
- ④ Check octet rule - each atom should have a share in 8 electrons (H gets 2). if not, make double or triple bonds.



In oxyacids, the acidic hydrogen atoms are attached to OXYGEN atoms in the structure!


$$\begin{array}{r} \text{H: } 1 \times 1 \\ \text{N: } 1 \times 5 \\ \text{O: } 2 \times 6 \\ \hline 18 e^- \end{array}$$



Here, the two oxygen atoms are in different chemical environments (the one on the right is bonded to both a nitrogen and a hydrogen atom), so we are not surprised to see different bonds form!

A DOT STRUCTURE FOR A LARGER MOLECULE

- ① Count valence electrons
- ② Pick central atom and draw skeletal structure
 - central atom is usually the one that needs to gain the most electrons!
 - skeletal structure has all atoms connected to center with single bonds
- ③ Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.
- ④ Check octet rule - each atom should have a share in 8 electrons (H gets 2). If not, make double or triple bonds.



$$\begin{array}{l|l} \text{C} : 4 \times 2 = 8 & \\ \text{H} : 1 \times 6 = 6 & \\ \text{O} : 6 \times 1 = 6 & \\ \hline & 20 \end{array}$$

This formula gives us clues about the structure of this large molecule. It has three parts, and three central atoms!

