

If 50.3 kJ of heat was released when 5.48 g of formic acid are burned at constant pressure, then what is the enthalpy change of this reaction per mole of formic acid?


$$Q = -50.3 \text{ kJ} ; \Delta H = \frac{Q_{\text{constant pressure}}}{\text{mol HCHO}_2}$$

Find moles formic acid:


$$5.48 \text{ g HCHO}_2 \times \frac{\text{mol HCHO}_2}{46.026 \text{ g HCHO}_2} = 0.1190631382 \text{ mol HCHO}_2$$

$$\Delta H = \frac{Q_{\text{constant pressure}}}{\text{mol HCHO}_2} = \frac{-50.3 \text{ kJ}}{0.1190631382 \text{ mol HCHO}_2} = \boxed{-422 \frac{\text{kJ}}{\text{mol HCHO}_2}}$$

Based on the calculation above, can we complete this thermochemical equation?



We calculated the enthalpy change at first PER MOLE OF FORMIC ACID (which is standard when doing a calorimetry experiment), but the reaction as written here uses TWO MOLES of formic acid. So, the enthalpy change that we put into this thermochemical equation is TWO TIMES what we calculated above.



What is the enthalpy change when 150. L of nitrogen monoxide are formed by this reaction at 25.0 C and 1.50 atm pressure?

1 - Convert 150 L NO to moles. Use IDEAL GAS EQUATION.

2 - Convert moles NO to enthalpy change using THERMOCHEMICAL EQUATION

$$PV = nRT \quad \left| \quad P = 1.50 \text{ atm} \quad R = 0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}} \right.$$

$$n = \frac{PV}{RT} \quad \left| \quad V = 150. \text{ L} \quad T = 25.0^\circ\text{C} = 298.2 \text{ K} \right.$$

$$\textcircled{1} \quad n_{\text{NO}} = \frac{(1.50 \text{ atm})(150. \text{ L})}{\left(0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}}\right)(298.2 \text{ K})} = 9.194822849 \text{ mol NO}$$

$$4 \text{ mol NO} = -906 \text{ kJ}$$

$$9.194822849 \text{ mol NO} \times \frac{-906 \text{ kJ}}{4 \text{ mol NO}} = \boxed{-2080 \text{ kJ}}$$

$\textcircled{2}$

34.086 g/mol  
-20.50

Heat of formation / enthalpy of formation!

0

-285.8

-296.8

$\Delta H_f^\circ$ , kJ/mol



(Appendix C)


What is the enthalpy change at standard conditions when 25.0 grams of hydrogen sulfide gas is reacted?

- 1 - Calculate the enthalpy change of the reaction as written. Use HESS'S LAW.
- 2 - Convert 25.0 grams hydrogen sulfide to moles. Use FORMULA WEIGHT.
- 3 - Convert moles hydrogen sulfide to enthalpy change. Use THERMOCHEMICAL EQUATION

$$\Delta H = \sum \Delta H_f^\circ, \text{ products} - \sum \Delta H_f^\circ, \text{ reactants}$$

$$= [2(-285.8) + 2(-296.8)] - [2(-20.50) + 3(0)] \quad (1)$$


$$= -1124.2 \text{ kJ}$$



$$34.086 \text{ g H}_2\text{S} = 1 \text{ mol H}_2\text{S} \quad | \quad 2 \text{ mol H}_2\text{S} = -1124.2 \text{ kJ}$$

$$25.0 \text{ g H}_2\text{S} \times \frac{1 \text{ mol H}_2\text{S}}{34.086 \text{ g H}_2\text{S}} \times \frac{-1124.2 \text{ kJ}}{2 \text{ mol H}_2\text{S}} = \boxed{-412 \text{ kJ}}$$

propane



Calculate the volume of propane gas at 25.0 C and 1.08 atm required to provide 565 kJ of heat using the reaction above.

- 1 - Convert the energy requirement to moles propane using THERMOCHEMICAL EQUATION.
- 2 - Convert moles propane to volume using IDEAL GAS EQUATION.

$$\text{mol C}_3\text{H}_8 = -2043 \text{ kJ}$$

Because the reaction (the system) is providing the energy, the energy requirement should be given a negative sign. The system gives up energy to the surroundings here...

$$-565 \text{ kJ} \times \frac{\text{mol C}_3\text{H}_8}{-2043 \text{ kJ}} = 0.2765540871 \text{ mol C}_3\text{H}_8 \quad (1)$$

$$PV = nRT$$

$$V = \frac{nRT}{P}$$

$$n = 0.2765540871 \text{ mol C}_3\text{H}_8$$

$$R = 0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}}$$

$$T = 25.0^\circ\text{C} = 298.2 \text{ K}$$



$$P = 1.08 \text{ atm}$$

$$V = \frac{(0.2765540871 \text{ mol C}_3\text{H}_8) \left(0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}}\right) (298.2 \text{ K})}{(1.08 \text{ atm})}$$

$$= 6.27 \text{ L C}_3\text{H}_8 @ 25.0^\circ\text{C}, 1.08 \text{ atm}$$

## END OF CHAPTER 6

## LIGHT



Diffraction

frequency = wavelengths / time =  $\nu$      $s^{-1}$ : Hertz, Hz

- Light has properties of WAVES such as DIFFRACTION (it bends around small obstructions).
- Einstein noted that viewing light as a particle that carried an energy proportional to the FREQUENCY could explain the PHOTOELECTRIC EFFECT!


$$E_{\text{photon}} = h \nu$$

Planck's constant:  $6.63 \times 10^{-34} \text{ J}\cdot\text{s}$

photon = particle or packet of light

(The photoelectric effect is the emission of electrons from a metal caused by exposure to light. Einstein discovered that if the light were not of the correct FREQUENCY, increasing the INTENSITY of the light would not cause electron emission. He concluded that individual photons must have enough energy to excite an electron - i.e. they must have the appropriate frequency.)

The photoelectric effect and Einstein's ideas about the energy content of light led us to discover a new model for the atom! How? Let's start with the nuclear model:



Nuclear model:

- Protons and neutrons in a dense NUCLEUS at center of atom
- Electrons in a diffuse (mostly empty) ELECTRON CLOUD surrounding NUCLEUS.

... so what's wrong with the nuclear model? Among other things, it doesn't explain ...