

A solution of 2.500g of unknown dissolved in 100.0 g of benzene has a freezing point of 4.880 C.

What is the molecular weight of the unknown?

$$K_{f, \text{benzene}} = 5.065 \text{ } ^\circ\text{C}/m, \quad T_{f, \text{benzene}} = 5.455 \text{ } ^\circ\text{C} \quad \left(\begin{array}{l} \text{see} \\ \text{p 500 9th} \\ \text{p 509, 10th} \end{array} \right)$$

$$\Delta T_f = K_f \times C_m \longrightarrow \frac{\text{mol unknown}}{\text{kg benzene}}$$

To find molecular weight, we need to find out how many moles of unknown we dissolved.

$$5.455 \text{ } ^\circ\text{C} - 4.880 \text{ } ^\circ\text{C} = 0.575 \text{ } ^\circ\text{C}$$

First, calculate the MOLAL CONCENTRATION of the solution!

$$0.575 \text{ } ^\circ\text{C} = (5.065 \text{ } ^\circ\text{C}/m) \times C_m$$

$$C_m = 0.1135241856 \text{ m} = \frac{0.1135241856 \text{ mol unkr}}{\text{kg benzene}}$$

To find MOLES UNKNOWN, multiply the MASS BENZENE by the MOLAL CONCENTRATION

$$0.1000 \text{ kg benzene} \times \frac{0.1135241856 \text{ mol unkr}}{\text{kg benzene}} = 0.0113524186 \text{ mol unkr}$$


MOLECULAR WEIGHT is mass per mole, so ...

$$MW = \frac{\text{mass unknown}}{\text{mol unknown}} = \frac{2.500 \text{ g}}{0.0113524186 \text{ mol unkr}} = \boxed{220. \text{ g/mol}}$$

Find C_m to find the moles of unknown!

75 VAPOR PRESSURE LOWERING

- Described by RAOULT'S LAW



P_A = partial pressure of the VAPOR of solvent molecules.

$$P_A = P_A^* \times X_A$$

mole fraction of component A

vapor pressure of pure component A (depends on temperature)

partial pressure of component A in a solution

... but component "A" above is actually the SOLVENT. If we want to describe this as a colligative property, we want to express Raolt's law in terms of the SOLUTE! Assuming a two-component mixture, we get...

$$\Delta P = P_A^* \times X_B$$

mole fraction of component B (the SOLUTE in a two-component mixture)

Vapor pressure lowering. This is the DECREASE in the vapor pressure of the solvent due to the presence of solute.

76 BOILING POINT ELEVATION

- Since the vapor pressure is lowered by the presence of a solute, AND since boiling occurs when the vapor pressure of a liquid equals the external pressure - solutes also cause BOILING POINT ELEVATION.

- The equation for boiling point elevation looks almost exactly like the equation for the freezing point depression, and is used in almost the same way.

$$\Delta T_b = K_b \times C_m$$

ΔT_b — Boiling point elevation: The amount the boiling temperature is RAISED by the solute.

K_b — Boiling point elevation constant (for SOLVENT) $\left(\begin{array}{l} \text{pSOc, 9th} \\ \text{pSO9, 10th} \end{array} \right)$

C_m — concentration of solute (molality)

What is the boiling point of a solution that contains 2.817 g of molecular sulfur (S_8) dissolved in 100.0 grams of acetic acid?

$$T_b = 118.5^\circ\text{C} \quad K_b = 3.08^\circ\text{C}/m \quad (\text{see p500 for data})$$

p509, 10th

$$\Delta T_b = K_b \times C_m \quad \left| \quad C_m = \frac{\text{mol } S_8}{\text{kg } \text{H}(\text{C}_2\text{H}_3\text{O}_2)} \right. \quad 0.1000 \text{ kg}$$

Find moles sulfur, then use that to find the MOLAL CONCENTRATION of the solution.

$$S_8: 8 \times 32.07 = 256.56 \text{ g } S_8 = \text{mol } S_8$$

$$2.817 \text{ g } S_8 \times \frac{\text{mol } S_8}{256.56 \text{ g } S_8} = 0.0109798877 \text{ mol } S_8$$

Find C_m :

$$C_m = \frac{0.0109798877 \text{ mol } S_8}{0.1000 \text{ kg } \text{H}(\text{C}_2\text{H}_3\text{O}_2)} = 0.1097988775 \text{ m}$$

Find BOILING POINT ELEVATION

$$\Delta T_b = K_b \times C_m = (3.08^\circ\text{C}/m)(0.1097988775 \text{ m}) = 0.3382^\circ\text{C}$$


Find new boiling point by adding ΔT to the original boiling point:

$$T_{b, \text{solution}} = 118.5^\circ\text{C} + 0.3382^\circ\text{C} = \boxed{118.8^\circ\text{C}}$$

OSMOTIC PRESSURE

permits flow of solvent, but not solute particles

- OSMOSIS: the flow of solvent molecules through a SEMIPERMEABLE membrane to equalize concentration of solute on each side of the membrane.



The rate of solvent migration towards the RIGHT is greater than that towards the LEFT.

If you apply enough pressure to the piston, osmosis will not occur. This pressure is called the OSMOTIC PRESSURE

$$\pi = M \times R \times T$$


temperature

ideal gas constant


molar concentration of solute

IONIC COMPOUNDS and colligative properties

- Ionic compounds DISSOCIATE in water into their component ions. Each ion formed can act as a solute and influence the colligative properties!



... so the concentration of IONS here is TWICE the nominal NaCl concentration.



... so the concentration of IONS here is THREE TIMES the nominal calcium chloride concentration.

- Ions interact with each other in solution, so unless an ionic solution is DILUTE, the effective concentrations of ions in solution will be less than expected. A more advanced theory (Debye-Huckel) covers this, but we'll assume that our solutions are dilute enough so that we can use the concentration of the ions in solution to determine the colligative properties!

If you are at an altitude high enough for the boiling point of water to be 95.00 C, what amount of sodium chloride would you need to add to 1.000 kg of water to raise the boiling point to 100.00 C?

$$K_b = 0.512 \text{ } ^\circ\text{C}/m \quad \text{NaCl: } 58.443 \text{ g/mol}$$

$$\Delta T_b = K_b \times C_m \quad \left| \quad C_m = \frac{\text{mol ions}}{\text{kg H}_2\text{O}} \right] 1.000 \text{ kg}$$


$$\left[\begin{array}{l} 100.00^\circ\text{C} - 95.00^\circ\text{C} = 5.00^\circ\text{C} \\ 0.512^\circ\text{C}/m \end{array} \right.$$

Find C_m (molal concentration of IONS)

$$5.00^\circ\text{C} = 0.512^\circ\text{C}/m \times C_m; \quad C_m = 9.765625 \text{ } m \text{ ions}$$

Find moles IONS ...

$$1.000 \text{ kg water} \times \frac{9.765625 \text{ mol ions}}{\text{kg water}} = 9.765625 \text{ mol ions}$$



Find moles sodium chloride, then convert to mass for weighing.

$$9.765625 \text{ mol ions} \times \frac{\text{mol NaCl}}{2 \text{ mol ions}} \times \frac{58.443 \text{ g NaCl}}{\text{mol NaCl}} = \boxed{285 \text{ g NaCl}}$$