


Example: How many grams of ammonium carbonate do we need to weigh out to get 3.65 moles of ammonium carbonate?

First, find the CHEMICAL FORMULA of ammonium carbonate



Now, find formula weight:

$$\text{N} : 2 \times 14.01$$

$$\text{H} : 8 \times 1.008$$

$$\text{C} : 1 \times 12.01$$

$$\text{O} : 3 \times 16.00$$

---


$$96.094 \text{ g } (\text{NH}_4)_2\text{CO}_3 = \text{mol } (\text{NH}_4)_2\text{CO}_3$$

Finally, calculate mass:

$$3.65 \text{ mol } (\text{NH}_4)_2\text{CO}_3 \times \frac{96.094 \text{ g } (\text{NH}_4)_2\text{CO}_3}{\text{mol } (\text{NH}_4)_2\text{CO}_3} = 351 \text{ g } (\text{NH}_4)_2\text{CO}_3$$

## PERCENTAGE COMPOSITION

- sometimes called "percent composition" or "percent composition by mass"
- the percentage of each element in a compound, expressed in terms of mass

Example: Find the percentage composition of ammonium nitrate.

$$\text{NH}_4\text{NO}_3 : \text{N} : 2 \times 14.01 = 28.02$$

$$\text{H} : 4 \times 1.008 = 4.032$$

$$\text{O} : 3 \times 16.00 = 48.00$$

$$\underline{80.052 \text{ g NH}_4\text{NO}_3 = 1 \text{ mol NH}_4\text{NO}_3}$$

These numbers are the masses of each element in a mole of the compound!

$$\% \text{ N} = \frac{28.02 \text{ g N}}{80.052 \text{ g total}} \times 100\% = 35.00\% \text{ N}$$

$$\% \text{ H} = \frac{4.032 \text{ g H}}{80.052 \text{ g total}} \times 100\% = 5.04\% \text{ H}$$


$$\% \text{ O} = \frac{48.00 \text{ g O}}{80.052 \text{ g total}} \times 100\% = 59.96\% \text{ O}$$

Check ... these should sum to 100% (within roundoff error)

So far, we have

- looked at how to determine the composition by mass of a compound from a formula
- converted from MASS to MOLES (related to the number of atoms/molecules)
- converted from MOLES to MASS

Use formula weight



Are we missing anything?

- What about SOLUTIONS, where the desired chemical is not PURE, but found DISSOLVED IN WATER?
- How do we deal with finding the moles of a desired chemical when it's in solution?

# MOLAR CONCENTRATION \*

- unit: MOLARITY (M): moles of dissolved substance per LITER of solution

$$M = \text{molarity} = \frac{\text{moles of SOLUTE}}{\text{L SOLUTION}}$$

↖ dissolved substance

$$6.0 \text{ M HCl solution} = \frac{6.0 \text{ mol HCl}}{\text{L}}$$

If you have 0.250 L (250 mL) of 6.0 M HCl, how many moles of HCl do you have?

$$6.0 \text{ mol HCl} = \text{L}$$

$$0.250 \text{ L} \times \frac{6.0 \text{ mol HCl}}{\text{L}} = 1.5 \text{ mol HCl}$$

\* See SECTIONS 4.7 - 4.10 for more information about MOLARITY and solution calculations (p 154 - 162 - 9th edition) (p 156-164 - 10th edition)

If you need 0.657 moles of hydrochloric acid, how many liters of 0.0555 M HCl do you need to measure out?

$$0.0555 \text{ mol HCl} = \text{L}$$

$$0.657 \text{ mol HCl} \times \frac{\text{L}}{0.0555 \text{ mol HCl}} = \boxed{11.8 \text{ L}}$$

11800 mL

This volume is much too large for typical lab-scale work!

We should use a more concentrated solution!

What if we used 6.00 M HCl?

$$6.00 \text{ mol HCl} = \text{L}$$

$$0.657 \text{ mol HCl} \times \frac{\text{L}}{6.00 \text{ mol HCl}} = \boxed{0.110 \text{ L}}$$


110 mL

This volume is more practical ... can be easily measured with a 250 mL cylinder.

Example: How would we prepare 500. mL of 0.500 M sodium sulfate in water?



Dissolve the appropriate amount of sodium sulfate into enough water to make 500. mL of solution.



A VOLUMETRIC FLASK is a flask that is designed to precisely contain a certain volume of liquid.

VOLUMETRIC FLASKS are used to prepare solutions.

volumetric flask

We know we need 500. mL solution, and we also know that solution should be 0.500 M. From that, we can calculate the moles of sodium sulfate. Then, we can convert the moles sodium sulfate to mass with the formula weight.

$$0.500 \text{ mol Na}_2\text{SO}_4 = \text{L} \quad | \quad \text{mL} = 10^{-3} \text{ L} \quad | \quad 142.05 \text{ g Na}_2\text{SO}_4 = \text{mol Na}_2\text{SO}_4$$

$$500. \text{ mL} \times \frac{10^{-3} \text{ L}}{\text{mL}} \times \frac{0.500 \text{ mol Na}_2\text{SO}_4}{\text{L}} \times \frac{142.05 \text{ g Na}_2\text{SO}_4}{\text{mol Na}_2\text{SO}_4} = \boxed{35.5 \text{ g Na}_2\text{SO}_4}$$

To prepare this solution, weigh 35.5 grams of sodium sulfate into a 500 mL volumetric flask, then dilute to the mark with distilled water.

## More on MOLARITY

To prepare a solution of a given molarity, you generally have two options:

① Weigh out the appropriate amount of solute, then dilute to the desired volume with solvent (usually water)

② Take a previously prepared solution of known concentration and DILUTE it with solvent to form a new solution

"stock solution"

- Use DILUTION EQUATION

The dilution equation is easy to derive with simple algebra.

$$M \times V$$

$$\frac{\text{mol}}{\text{L}} \times \text{L} = \text{moles solute}$$

... but when you dilute a solution, the number of moles of solute REMAINS CONSTANT. (After all, you're adding only SOLVENT)

$$M_1 V_1 = M_2 V_2$$

before  
dilution

after  
dilution

Since the number of moles of solute stays the same, this equality must be true!

$$M_1 V_1 = M_2 V_2 \quad \dots \text{the "DILUTION EQUATION"}$$

$M_1$  = molarity of concentrated solution

$V_1$  = volume of concentrated solution

$M_2$  = molarity of dilute solution

$V_2$  = volume of dilute solution (total volume, not volume of added solvent!)

The volumes don't HAVE to be in liters, as long as you use the same volume UNIT for both volumes!

Example: Take the 0.500 M sodium sulfate we discussed in the previous example and dilute it to make 150. mL of 0.333 M solution. How many mL of the original solution will we need to dilute?

$$M_1 V_1 = M_2 V_2 \quad M_1 = 0.500 \text{ M} \quad M_2 = 0.333 \text{ M}$$

$$V_1 = ??? \quad V_2 = 150. \text{ mL}$$

$$(0.500 \text{ M}) V_1 = (0.333 \text{ M})(150. \text{ mL})$$


$$V_1 = \boxed{99.9 \text{ mL of } 0.500 \text{ M Na}_2\text{SO}_4}$$

Take 99.9 mL of 0.500 M sodium sulfate, then add enough distilled water to make 150. mL of solution. (Ideally, transfer the 0.500 M sodium sulfate to a 150 mL volumetric flask and dilute to the mark with distilled water...)



## CHEMICAL CALCULATIONS CONTINUED: REACTIONS

- Chemical reactions proceed on an ATOMIC basis, NOT a mass basis!
- To calculate with chemical reactions (i.e. use chemical equations), we need everything in terms of ATOMS ... which means MOLES of atoms



coefficients are in terms of atoms and molecules!



- To do chemical calculations, we need to:
  - Relate the amount of substance we know (mass or volume) to a number of moles
  - Relate the moles of one substance to the moles of another using the equation
  - Convert the moles of the new substance to mass or volume as desired