



THE KINETIC PICTURE OF GASES



① Gas molecules are small compared to the space between the gas molecules!

LOW DENSITY!



- ② Gas molecules are constantly in motion. They move in straight lines in random directions and with various speeds.
- ③ Attractive and repulsive forces between gas molecules are so small that they can be neglected except in a collision.
 - Each gas molecule behaves independently of the others.
- ④ Collisions between gas molecules and each other or the walls are ELASTIC.

⑤ The average kinetic energy of gas molecules is proportional to the absolute temperature.

How does this picture explain the properties of gases?

- Gases expanding to fill their container? Agrees with kinetic picture, since gas molecules are independent
- Thermal expansion of gas at constant pressure? Agrees, because the container has to EXPAND to keep the pressure (from collisions) constant when the gas molecules move faster.
- Pressure increases with temperature at constant volume: Agrees, because the number and force of collisions increases with molecular speed.

GAS LAWS

- were derived by experiment long before kinetic theory, but agree with the kinetic picture!

Boyle's Law:

$$PV = \text{constant} \quad \left. \vphantom{PV = \text{constant}} \right] \text{ True at constant temperature}$$

$$P_1 V_1 = \text{constant}$$

$$P_2 V_2 = \text{constant}$$

$$\left. \vphantom{P_1 V_1 = \text{constant}} \right] \rightarrow \boxed{P_1 V_1 = P_2 V_2} \quad \text{True at constant temperature}$$

Charles's Law:

$$\frac{V}{T} = \text{constant} \quad \left. \vphantom{\frac{V}{T} = \text{constant}} \right] \text{ True at constant pressure, and using ABSOLUTE temperature}$$

$$\left. \vphantom{\frac{V}{T} = \text{constant}} \right] \rightarrow \boxed{\frac{V_1}{T_1} = \frac{V_2}{T_2}} \quad \text{True at constant pressure, and using ABSOLUTE temperature}$$

140 Combined gas law:

$$\frac{PV}{T} = \text{constant}$$

Must use ABSOLUTE temperature units!

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

Must use ABSOLUTE temperature units!


↑ amount (moles) of gas must be constant!

Avogadro's law:

- a mole of any gas at the same conditions has the same volume.

1 mol gas molecules @ 0°C and 1 atm
volume = 22.4 L

"STP"
Standard
Temperature
and
Pressure



Ideal gas law:

$$\frac{PV}{T} = \text{constant}$$

... but this constant actually depends on the amount of gas!

$$= n \times "R"$$

The ideal gas constant,

$$0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}}$$

... combining these together ...

$$\frac{PV}{T} = nR$$



$$PV = nRT$$

P = pressure atm

V = volume L

T = ABSOLUTE temperature K

R = ideal gas constant

n = number of moles of gas molecules

A balloon is taken from a room where the temperature is 27.0 C to a freezer where the temperature is -5.0 C. If the balloon has a volume of 3.5 L in the 27.0 C room, what is the volume of the balloon in the freezer. Assume pressure is constant.

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}, \text{ with } P \text{ constant}, \quad \frac{V_1}{T_1} = \frac{V_2}{T_2}$$

$$V_1 = 3.5 \text{ L}$$

$$T_1 = 27.0^\circ\text{C} = 300.2 \text{ K} \quad \left| \quad \begin{array}{l} V_2 = ? \\ T_2 = -5.0^\circ\text{C} = 268.2 \text{ K} \end{array} \right.$$

$$\frac{3.5 \text{ L}}{300.2 \text{ K}} = \frac{V_2}{268.2 \text{ K}}; \quad V_2 = \boxed{3.1 \text{ L in freezer}}$$

2.25 L of nitrogen gas is trapped in a piston at 25.0 C and 1.00 atm pressure. If the piston is pushed in so that the gas's volume is 1.00 L while the temperature increases to 31.0 C, what is the pressure of the gas in the piston?

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} \quad \begin{array}{l} P_1 = 1.00 \text{ atm} \\ V_1 = 2.25 \text{ L} \\ T_1 = 25.0^\circ\text{C} = 298.2 \text{ K} \end{array} \quad \left| \quad \begin{array}{l} P_2 = ? \\ V_2 = 1.00 \text{ L} \\ T_2 = 31.0^\circ\text{C} = 304.2 \text{ K} \end{array} \right.$$

$$\frac{(1.00 \text{ atm})(2.25 \text{ L})}{(298.2 \text{ K})} = \frac{P_2 (1.00 \text{ L})}{(304.2 \text{ K})}; \quad P_2 = \boxed{2.30 \text{ atm}}$$

Calculate the mass of 22650 L^{*} of oxygen gas at 25.0 C and 1.18 atm pressure.



* Volume of a 10'x10'x8' room

Use the ideal gas equation to calculate MOLES of oxygen gas. Then, use FORMULA WEIGHT of oxygen gas to find the mass.


$$PV = nRT$$

$$n = \frac{PV}{RT} \quad \left| \begin{array}{l} P = 1.18 \text{ atm} \\ V = 22650 \text{ L} \\ T = 25.0^\circ\text{C} = 298.2 \text{ K} \end{array} \right. \quad \begin{array}{l} n = ? \\ R = 0.08206 \frac{\text{L}\cdot\text{atm}}{\text{mol}\cdot\text{K}} \end{array}$$

$$n_{\text{O}_2} = \frac{(1.18 \text{ atm})(22650 \text{ L})}{\left(0.08206 \frac{\text{L}\cdot\text{atm}}{\text{mol}\cdot\text{K}}\right)(298.2 \text{ K})} = 1092.22235 \text{ mol O}_2$$

$$1092.22235 \text{ mol O}_2 \times \frac{32.00 \text{ g O}_2}{\text{mol O}_2} = \boxed{35000 \text{ g O}_2} \quad \begin{array}{l} 35.0 \text{ kg} \\ \sim 77 \text{ lb} \end{array}$$

$$FW_{\text{NaHCO}_3} = 84.007 \text{ g/mol}$$



Given 25.0 g of sodium bicarbonate and sufficient sulfuric acid, what volume of carbon dioxide gas would be produced at 25.0 C and 0.950 atm pressure?

- 1 - Convert 25.0 g sodium bicarbonate to moles using FORMULA WEIGHT.
- 2 - Convert moles sodium bicarbonate to moles carbon dioxide gas using CHEMICAL EQUATION
- 3 - Convert moles carbon dioxide gas to volume using IDEAL GAS EQUATION.

$$84.007 \text{ g NaHCO}_3 = 1 \text{ mol NaHCO}_3 \quad | \quad 2 \text{ mol NaHCO}_3 = 2 \text{ mol CO}_2$$

$$25.0 \text{ g NaHCO}_3 \times \frac{1 \text{ mol NaHCO}_3}{84.007 \text{ g NaHCO}_3} \times \frac{2 \text{ mol CO}_2}{2 \text{ mol NaHCO}_3} = 0.2975942461 \text{ mol CO}_2$$

$PV = nRT$	$n = 0.2975942461 \text{ mol CO}_2$	$R = 0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}}$
$V = \frac{nRT}{P}$	$T = 25.0^\circ\text{C} = 298.2 \text{ K}$	
	$P = 0.950 \text{ atm}$	

$$V = \frac{(0.2975942461 \text{ mol CO}_2) \left(0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}}\right) (298.2 \text{ K})}{(0.950 \text{ atm})} = \boxed{7.67 \text{ L}}$$

... at 25.0 C
and 0.950 atm

What volume would the gas in the last example problem have at STP?

STP: "Standard Temperature and Pressure" (0 C and 1 atm)

Let's solve this using the combined gas law:

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} \quad \left| \quad \begin{array}{l} P_1 = 0.950 \text{ atm} \\ V_1 = 7.665492788 \text{ L} \\ T_1 = 298.2 \text{ K} \end{array} \quad \begin{array}{l} P_2 = 1 \text{ atm} \\ V_2 = ? \\ T_2 = 273.2 \text{ K} \end{array} \right.$$

$$\frac{(0.950 \text{ atm})(7.665492788 \text{ L})}{(298.2 \text{ K})} = \frac{(1 \text{ atm})(V_2)}{(273.2 \text{ K})}$$

$$V_2 = \boxed{6.67 \text{ L}} \text{ at STP}$$

Alternate solution: Use the ideal gas equation

$$PV = nRT \quad \left| \quad \begin{array}{l} n = 0.2975942481 \text{ mol CO}_2 \\ R = 0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}} \end{array} \right.$$

$$V = \frac{nRT}{P} \quad \left| \quad \begin{array}{l} T = 273.2 \text{ K} \\ P = 1 \text{ atm} \end{array} \right.$$

$$V = \frac{(0.2975942481 \text{ mol CO}_2) \left(0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}}\right) (273.2 \text{ K})}{(1 \text{ atm})}$$

$$= 6.67 \text{ L at STP}$$



At 300°C , ammonium nitrate violently decomposes to produce nitrogen gas, oxygen gas, and water vapor. What is the total volume of gas that would be produced at 1.00 atm by the decomposition of 15.0 grams of ammonium nitrate?

To simplify this problem, let's calculate TOTAL MOLES OF GAS instead of the individual moles of each gas.

$$F_w \text{NH}_4\text{NO}_3 \approx 80.052 \text{ g/mol}$$

- 1 - Convert 15.0 grams ammonium nitrate to moles using FORMULA WEIGHT
- 2 - Convert moles ammonium nitrate to TOTAL MOLES OF GAS using chemical equation.
- 3 - Convert TOTAL MOLES OF GAS to volume using IDEAL GAS EQUATION.

$$80.052 \text{ g NH}_4\text{NO}_3 = 1 \text{ mol NH}_4\text{NO}_3 \quad | \quad 2 \text{ mol NH}_4\text{NO}_3 = 7 \text{ mol gas } (2+1+4)$$

$$15.0 \text{ g NH}_4\text{NO}_3 \times \frac{1 \text{ mol NH}_4\text{NO}_3}{80.052 \text{ g NH}_4\text{NO}_3} \times \frac{7 \text{ mol gas}}{2 \text{ mol NH}_4\text{NO}_3} = 0.6558237146 \text{ mol gas}$$

$$PV = nRT \quad | \quad n = 0.6558237146 \text{ mol gas} \quad R = 0.08206 \frac{\text{L}\cdot\text{atm}}{\text{mol}\cdot\text{K}}$$

$$V = \frac{nRT}{P} \quad | \quad T = 300.^\circ\text{C} = 573 \text{ K} \quad P = 1.00 \text{ atm}$$

$$V = \frac{(0.6558237146 \text{ mol gas}) (0.08206 \frac{\text{L}\cdot\text{atm}}{\text{mol}\cdot\text{K}}) (573 \text{ K})}{(1.00 \text{ atm})} = \boxed{30.8 \text{ L}}$$