


... but using electron configurations to describe how aluminum bromide forms is a bit cumbersome! Can we simplify the picture a bit?

## LEWIS NOTATION / ELECTRON-DOT NOTATION

- Lewis notation represents each VALENCE electron with a DOT drawn around the atomic symbol. Since the valence shell of an atom contains only "s" and "p" electrons, the maximum number of dots drawn will be EIGHT.

- To use electron-dot notation, put a dot for each valence electron around the atomic symbol. Put one dot on each "side" of the symbol (4 sides), then pair the dots for atoms that have more than four valence electrons.

examples:



More examples



Which "side" you draw the dots on isn't important, as long as you have the right number of electrons and the right number of "pairs"



To draw a dot structure for an atom, you need to know HOW MANY valence electrons it has! You can determine this simply from the periodic table, WITHOUT writing the whole electron configuration!

The number of valence electrons equals the group number in the A/B group numbering system FOR "A" GROUPS!

	IA												VIII A								
1	H	IIA											III A	IVA	VA	VIA	VIIA	He			
2	Li	Be											B	C	N	O	F	Ne			
3	Na	Mg	IIIB	IVB	VB	VIB	VIIB	VIIIB	IB	IIB						Al	Si	P	S	Cl	Ar
4	K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr			
5	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe			
6	Cs	Ba	La*	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn			
7	Fr	Ra	Ac*	Rf	Db	Sg	Bh	Hs	Mt												

2 valence electrons

1 valence electron

3 valence electrons

4 valence electrons


5 valence electrons

6 valence electrons

7 valence electrons

8 valence electrons (except helium!)

... but how do we use this to describe a reaction that produces ions? Let's look at our previous example!




... this is a bit easier to follow than looking at all those letters and numbers in the electron configurations for these elements!

## MOLECULAR COMPOUNDS


- Form when atoms SHARE electrons instead of transferring them. This results in the formation of MOLECULES ... groups of atoms held together by electron-sharing.

How might atoms SHARE electrons? By coming together close enough so that their atomic ORBITALS overlap each other:

This idea is called  
VALENCE  
BOND  
THEORY!



Each hydrogen atom has a single electron in a 1s orbital.




When hydrogen atoms come close enough to each other for these orbitals to OVERLAP, each hydrogen "sees" BOTH electrons, filling up the "s" orbitals of both atoms. This is a COVALENT BOND.

Co: together  
Valent: valence shells

This is the DIATOMIC MOLECULE,  $H_2$

... so how would this look using dot notation?




In dot structures, SHARED PAIRS of electrons are often written as DASHES to make the structures look neater.



\* Why doesn't hydrogen end up with eight electrons? Because hydrogen has only the first shell, which contains only a single "s" subshell (NO "p" subshell). This "s" subshell is full with two electrons, and that's all hydrogen needs to get.

Let's look at OXYGEN ...



We know that oxygen exists in air as the diatomic molecule  $O_2$



The oxygen atoms share TWO pairs of electrons. This is called a DOUBLE BOND

OR




Each oxygen atom has a share in eight electrons!


A few notes on the double bond:

- ① - For atoms to share more than one pair of electrons, they have to move closer to one another than they would if they were only sharing one pair of electrons. This BOND DISTANCE is measurable!
- ② - It takes more energy to break a double bond between two atoms than it would to break a single bond between the same two atoms. This BOND ENERGY is also measurable!

Let's look at NITROGEN ...



We know that nitrogen exists in air as the diatomic molecule  $N_2$



The nitrogen atoms share THREE pairs of electrons. This is called a TRIPLE BOND

OR



Nitrogen gas is fairly inert ... it's hard to break the triple bond in nitrogen gas apart!

A few notes on the triple bond:

- ① - For atoms to share three pairs of electrons, they have to move closer to one another than they would if they were sharing one or two pairs of electrons. Triple bonds have the shortest BOND DISTANCE of all covalent bonds.
- ② - It takes more energy to break a triple bond between two atoms than it would to break either a single or double bond between the same two atoms. The triple bond has the largest BOND ENERGY of all three kinds of covalent bonds.

SO FAR, we've seen that ...





- ① Atoms may share one, two, or three pairs of electrons with each other.
- ② Atoms will usually share enough electrons so that each atom ends up with a share in EIGHT electrons - the "octet rule"

- HYDROGEN will only end up with two electrons!

- Some other atoms may end up with more or less than eight electrons. Exceptions to the octet rule are covered in Chapter 9.

NOW, how could we come up with dot structures for some more complicated (and therefore, more interesting) molecules?

Examples:





① Count valence electrons

② Pick central atom and draw skeletal structure

- central atom is usually the one that needs to gain the most electrons!

- skeletal structure has all atoms connected to center with single bonds

③ Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.

④ Check octet rule - each atom should have a share in 8 electrons (H gets 2). if not, make double or triple bonds.




$$\text{C} : 1 \times 4$$

$$\text{O} : 1 \times 6$$

$$\text{Cl} : 2 \times 7 = 14$$

---



$$24e^-$$



Choose CARBON as the central atom since it needs to gain more electrons than the others...




Distribute remaining electrons, stop when we run out.



... but the central carbon atom has a share in only SIX electrons.!

To get carbon more electrons, we'll need to make a DOUBLE BOND. But with which other atom? Pick OXYGEN here, since it needed more electrons than chlorine did...




This structure looks better, since all atoms have a share in eight electrons!

- Count valence electrons
- Pick central atom and draw skeletal structure

- central atom is usually the one that needs to gain the most electrons!
- skeletal structure has all atoms connected to center with single bonds

- Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.

- Check octet rule - each atom should have a share in 8 electrons (H gets 2). If not, make double or triple bonds.




---


$$18 e^-$$



We will pick NITROGEN for the central atom, since it needs more electrons than oxygen or chlorine.



We ran out of space on the outer atoms, so the last two electrons go on the central nitrogen

Even with this pair of electrons on the central nitrogen atom, there still aren't enough. So we'll need a double bond. Just like last time, we'll make the double bond with OXYGEN:



Now all atoms have a share in eight electrons!


① Count valence electrons

② Pick central atom and draw skeletal structure


- central atom is usually the one that needs to gain the most electrons!
- skeletal structure has all atoms connected to center with single bonds

③ Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.


④ Check octet rule - each atom should have a share in 8 electrons (H gets 2). If not, make double or triple bonds.




Pick carbon as central atom and distribute electrons...



Carbon has a share in four electrons...



... now six.



Adding a second double bond gives carbon a share in eight electrons!



What's wrong with the green structure? The two oxygen atoms are in the same environment and SHOULD bond the same way!

If this were the correct structure, we'd see two different BOND LENGTHS in carbon dioxide ... but experimentally we see only one ... suggesting the double-bond structure we drew earlier was correct.

① Count valence electrons

② Pick central atom and draw skeletal structure

- central atom is usually the one that needs to gain the most electrons!


- skeletal structure has all atoms connected to center with single bonds

③ Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.


④ Check octet rule - each atom should have a share in 8 electrons (H gets 2). If not, make double or triple bonds.




In oxyacids, the acidic hydrogen atoms are attached to OXYGEN atoms in the structure!



18e<sup>-</sup>



OXYACID, so we must have a hydrogen atom attached to an oxygen atom!



... but NITROGEN has a share in only six electrons!



Unlike the carbon dioxide structure, these two oxygen atoms are in DIFFERENT environments, and bond to the central nitrogen differently.

## A DOT STRUCTURE FOR A LARGER MOLECULE

$$\begin{array}{l}
 \text{C} : 4 \times 2 = 8 \\
 \text{H} : 1 \times 6 = 6 \\
 \text{O} : 6 \times 1 = 6
 \end{array}
 \left| \begin{array}{l} \\ \\ \\ \hline 20
 \end{array} \right.$$


① Count valence electrons

② Pick central atom and draw skeletal structure

- central atom is usually the one that needs to gain the most electrons!
- skeletal structure has all atoms connected to center with single bonds

③ Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.

④ Check octet rule - each atom should have a share in 8 electrons (H gets 2). If not, make double or triple bonds.



This formula gives us a hint to the structure of ethanol. Ethanol has THREE central atoms chained together.

