

- What good is ionic theory?

- provides an easy-to-understand MECHANISM for certain kinds of chemical reactions.

- "Exchange" reactions. (a.k.a "double replacement" reactions)



Looking a bit more closely...



$$A_{g}^{+}(a_{q}) + (1^{-}(a_{q}) \rightarrow A_{g}C(s)]$$
 "Net ionic equation"

(The net ionic equation shows only ions and substances that change during the course of the reaction!)

- The net ionic equation tells us that any source of aqueous silver and chloride ions will exhibit this same chemistry, not just silver nitrate and sodium chloride!

<sup>115</sup> A bit more about molecular, ionic, and net ionic equations

- molecular equations: Represent all substances (even ionic substances) as if they were molecules. Include spectator ions, and do not show charges on ions. Traditional chemical equations.

- ionic equations: Show all free ions - including spectators - in a chemical reaction. Molecules and WEAK electrolytes are shown as molecules. STRONG electrolytes (like HCI) are shown as ions. Ions that are part of <u>undissolved ionic compounds</u> are shown as molecules.

- NET ionic equation: An ionic equation that leaves out spectator ions. Intended to show only things that actually change in a reaction.

$$\begin{array}{l} \operatorname{Ag}\operatorname{NO}_{2}(\operatorname{aq}) + \operatorname{Nu}\operatorname{Cl}(\operatorname{au}) \xrightarrow{\rightarrow} \operatorname{Ag}\operatorname{Cl}(\operatorname{s}) + \operatorname{Nu}\operatorname{No}_{2}(\operatorname{aq}) \\ \operatorname{Ag}^{\dagger}(\operatorname{au}) + \operatorname{No}_{2}^{-}(\operatorname{au}) + \operatorname{Na}^{\dagger}(\operatorname{au}) + \operatorname{Cl}^{-}(\operatorname{au}) \xrightarrow{\rightarrow} \operatorname{Ag}\operatorname{Cl}(\operatorname{s}) + \operatorname{Na}^{\dagger}(\operatorname{au}) + \operatorname{No}_{2}^{-}(\operatorname{au}) \\ \operatorname{Ag}^{\dagger}(\operatorname{au}) + \operatorname{Cl}^{-}(\operatorname{au}) \xrightarrow{\rightarrow} \operatorname{Ag}\operatorname{Cl}(\operatorname{s}) \end{array}$$

\* You can get from the complete ionic equation to the net ionic equation by crossing out the spectator ions on both sides.

## "Undissolved ionic compounds":

How can I tell if an ionic compound dissolves in water?

- consult experimental data: "solubility rules"!
  - A few of the "rules"...
    - Compounds that contain a Group IA cation (or ammonium) are soluble
    - Nitrates and acetates are soluble
    - Carbonates, phosphates, and hydroxides tend to be insoluble

... or see the web site for a solubility chart.

Fe(OH)3

#8 - hydroxides generally insoiluble, except Group IA, ammonium, calcium strontium, barium

Conclusion: iron(III) hydroxide is insoluble.

Hg L #3 - lodides usually dissolve, exceptions are silver, mercury, lead

Conclusion: silver(I) iodide is INSOLUBLE

$$Ca(C_2H_3O_2)_2$$

#2 - acetates are soluble, no common exceptions.

Conclusion: calcium acetate is soluble.




#5 - Most carbonates are insoluble

Conclusion - barium carbonate is insoluble.

Exchange Chemistry

- Three kinds of exchange chemistry.



Just because you mix together two ionic compounds does NOT mean that a reaction will occur. You need a DRIVING FORCE for a reaction.

118

#### PRECIPITATION REACTIONS

- driving force is the formation of an insoluble ionic compound.

Maz  $(POy)_{\gamma}(s) + 6 N \alpha C | (aq)$  $3Mg(l_2(aq) + 2Na_3POy(aq) \rightarrow$ ions: N Nat P04 3-The formation of SOLID (insoluble) magnesium phosphate is the driving force for this reaction.

When you're trying to complete a precipitation reaction:

(i) Write the IONS that form when the reactants are dissolved.

Description Make NEW compounds by pairing up cations with anions. Don't forget that the positive and negative charges must balance each other out!

(3) Use the solubility rules to determine the PHASE of each new compound - solid or aqueous.

(4)Balance the overall equation.

$$M_{g}(I_{2}(a_{q}) + N_{a}(_{2}H_{3}O_{2}(a_{q}) \rightarrow NO \text{ REACTION!}^{*})$$
ions:  $M_{g}^{2+} CI^{-} N_{a}^{+} (_{2}H_{3}O_{2}^{-})$ 

$$(exchange)^{*}$$

$$M_{g}((_{2}H_{3}O_{2})_{2} \dots \text{ dissolves in water})$$

$$N_{a}(I_{a} \dots \text{ dissolves in water})$$

So, no solid forms here. All possible combinations of these four ions result in compounds that dissolve readily in water.

$$\frac{m_g^{2+}Cl^{-}}{\log^2 Cl^{-}} + \frac{N_0^{+}C_2H_3U_2^{-}}{N_0^{+}} \rightarrow \frac{N_0^{+}m_g^{2+}}{\log^2 Cl^{-}}$$

$$\frac{N_0^{+}M_g^{2+}C_2H_3U_2^{-}}{\log^2 Cl^{-}}$$


NO CHANGE, therefore NO DRIVING FORCE, and NO REACTION

★ We will learn about other driving forces than the formation of solid, but these driving forces do not apply to this reaction

# ACID/BASE REACTIONS (also called NEUTRALIZATION REACTIONS)

- There are several stable molecules that may be formed in double replacement reactions, but the most common is <u>WATER</u>!

- Double replacement reactions that form water are also called "neutralizations"



\* To make water (  $H_2O$  ), you need a source of hydrogen ion (  $H^4$  ) and hydroxide ion (  $GH^-$  )

$$H^{+}(aq) + OH^{-}(aq) \longrightarrow H_{2}O(\ell)$$

$$\begin{array}{c} \text{This is the} \\ \text{NET IONIC} \\ \text{EQUATION} \\ \text{for many} \\ \text{neutralizations} \\ \\ \text{STRONG base!} \end{array}$$

### ACIDS

- compounds that release hydrogen ion (H<sup>+</sup>), when dissolved in water. Properties of acids:
  - Corrosive: React with most metals to give off hydrogen gas
  - Cause chemical burns on contact
  - Taste sour (like citrus citric acid!)
  - Changes litmus indicator to RED

### BASES

- Substances that release hydroxide ion (OH~) when dissolved in water

### Properties of bases:

- Caustic: Attack and dissolve organic matter (think lye, which is NaOH)
- Cause skin/eye damage on contact
- Taste bitter
- changes litmus indicator to BLUE

Due to the dissolving action of base on your skin, bases will feel "slippery". The base ITSELF is not particularly slippery, but what's left of your skin IS! <sup>122</sup> ACID/BASE or NEUTRALIZATION reactions continued

- the driving force of these reactions is the formation of water molecules.

$$\begin{array}{c} H^{+}(u_{q}) + OH^{-}(u_{q}) \longrightarrow H_{2}O(\theta) \end{array} \text{ Net ionic equation} \\ \hline H_{2}So_{y}(u_{q}) + 2N_{a}OH(u_{q}) \longrightarrow 2H_{2}O(\theta) + Na_{2}SO_{y}(u_{q}) \\ \hline H^{+}So_{y} & Na^{+}OH^{-} & The driving force for this reaction is the formation of water MOLECULES. \\ - How can this reaction be detected? \end{array}$$

- pH detector (indicator paper, etc.)
- do the products have similar chemical properties to the reactants?

- r<u>elease of he</u>at!

... formation of water is usually accompanied by a release of heat

### GAS FORMATION / OTHER MOLECULES

- There are a few other molecules that can be made with exchange-type chemistry.
- Most of these molecules are unstable and can break apart to form gases.
- Formation of a weak acid:
  - The formation of ANY weak acid in an exchange-type reaction can be a driving force.
  - Some weak acids are unstable and can break apart into gas molecules.

$$H_2(O_3Lag) \longrightarrow H_2O(l) + (CO_2C_g)$$
 Gas bubbles can leave solution!

... but how would you form carbonic acid in an exchange-type reaction?

acid + carbonate 
$$(0_3^{2-1})^{OR}$$
  
acid + bicarbonate  $H(0_3^{-1})^{OR}$   
 $H_2SO_4(a_4)+2N_4H(0_3(a_4)) \rightarrow 2H_2CO_3(a_4)+Na_2SO_4(a_4)$   
 $H^+SO_4^{1-1}N_a^+H(0_3^{-1})^{OR}$   
Formation of carbonic acid  
drives the reaction ... BUT ...

ł

... but when we mix sulfuric acid and sodium bicarbonate, we observe BUBBLES. We need to write an equation that agrees with our observations. We know that carbonic acid decomposes, so we go ahead and put that into our equation.

$$H_2(O_3Lag) \longrightarrow H_2O(l) + CO_2(g)$$

$$H_2O_3Lag) \rightarrow Na_2SO_4(ag) + 2H_2O(l) + 2(O_2(g))$$

Other molecules of interest:

$$\begin{array}{l} H_2 SO_3 : \text{ sulfurous acid } - \text{React an ACID with a SULFITE} \\ H_2 SO_3(n_q) \rightarrow H_2 O(l) + SO_2(g) \\ H_2 S : \text{ hydrogen sulfide (gas) - React an ACID with a SULFIDE} \\ H_2 SO_4(n_q) + N_{a_2} S(n_q) \rightarrow N_{a_2} So_4(n_q) + H_2 S(g) \end{array}$$