

Calculate how many grams of acrylonitrile could be obtained from 651000 g of propylene, assuming there is excess NO present.

- 1 - Convert mass propylene to moles propylene. Use FORMULA WEIGHT.
- 2 - Convert moles propylene to moles acrylonitrile. Use CHEMICAL EQUATION.
- 3 - Convert moles acrylonitrile to mass acrylonitrile. Use FORMULA WEIGHT.

$$\textcircled{1} \quad 42.081 \text{ g C}_3\text{H}_6 = \text{mol C}_3\text{H}_6$$

$$\textcircled{2} \quad 4 \text{ mol C}_3\text{H}_6 = 4 \text{ mol C}_3\text{H}_3\text{N}$$


$$\textcircled{3} \quad 53.064 \text{ g C}_3\text{H}_3\text{N} = \text{mol C}_3\text{H}_3\text{N}$$

$$651000 \text{ g C}_3\text{H}_6 \times \frac{\text{mol C}_3\text{H}_6}{42.081 \text{ g C}_3\text{H}_6} \times \frac{4 \text{ mol C}_3\text{H}_3\text{N}}{4 \text{ mol C}_3\text{H}_6} \times \frac{53.064 \text{ g C}_3\text{H}_3\text{N}}{\text{mol C}_3\text{H}_3\text{N}} =$$

$$= \boxed{821000 \text{ g C}_3\text{H}_3\text{N}}$$

## CONCEPT OF LIMITING REACTANT

- When does a chemical reaction STOP?




- When does this reaction stop? When burned in open air, this reaction stops when all the MAGNESIUM STRIP is gone. We say that the magnesium is LIMITING.

- This reaction is controlled by the amount of available magnesium

- At the end of a chemical reaction, the LIMITING REACTANT will be completely consumed, but there may be some amount of OTHER reactants remaining. We do chemical calculations in part to minimize these "leftovers".

- Reactants that are left at the end of a chemical reaction (in other words, they are NOT the limiting reactant) are often called "excess". So reacting magnesium with "excess oxygen" means that magnesium is limiting.

## STRUCTURE OF THE ELECTRON CLOUD



The nuclear model describes atoms as consisting of a NUCLEUS containing protons and neutrons and an ELECTRON CLOUD containing electrons.

The ELECTRON CLOUD is described as being a diffuse (lots of empty space) region of the atom. Nothing else about it is part of the nuclear model.


... but the nuclear model is not useful to explain several things:

- Does not explain why atoms react differently from one another
- Does not explain how atoms emit and absorb light (atomic line spectra)


# <sup>156</sup> ELECTROMAGNETIC SPECTRUM

(see p240-241)


- Different kinds of "light" have different energy contents




- Different colors of visible light correspond to different amounts of energy





Source of white light




Prism




Rainbow (all colors represented)

Gaseous Helium excited  
by electricity

Prism

LINE SPECTRUM - only  
a few specific colors appear!  
(see p<sup>243</sup> for example)

- Atomic line spectra are UNIQUE to each element. They're like atomic "fingerprints".
- Problem was that the current model of the atom completely failed to explain why atoms emitted these lines.




An orbit that is FARTHER from the nucleus means that the electron has MORE energy

An orbit that is CLOSER to the nucleus means that the electron has LESS energy

- Electrons may gain or lose energy by either ABSORBING (to gain) or EMITTING (to lose) a PHOTON of light. (Photon = particle or "packet" of energy.)
- If the electrons can gain or lose ANY amount of energy, then each atom would emit a RAINBOW rather than an LINE SPECTRUM.

## BOHR MODEL

- Theorized that electrons couldn't be just ANYWHERE around the nucleus. There must be restrictions on the motion of electrons that traditional physics did not explain.




- theorized that electrons could only be certain distances from the nucleus. In other words, they could only have certain values for ENERGY.

- Electrons could move only from one "energy level" to another DIRECTLY by giving up or absorbing a photon (light) that was equal in energy to the distance between the energy levels.

- The restrictions on where electrons could be in Bohr's model predicted that atoms would give LINE SPECTRA.

- Bohr's model accurately described the line spectrum of hydrogen (first time this had been done!)

- For other atoms, Bohr's model predicted a line spectrum, but the lines weren't the right colors!



Bohr's model didn't account for electron-electron interactions (which didn't exist in HYDROGEN)

- To account for this added complexity, a more sophisticated model had to be devised: QUANTUM THEORY. Quantum theory is the modern picture of the atom and its electron cloud.




## SHELLS, SUBSHELLS, AND ORBITALS

- Bohr's model predicted that energy levels (called SHELLS) were enough to describe completely how electrons were arranged around an atom. But there's more to it!


SHELL: Equivalent to Bohr's energy levels. Electrons in the same SHELL are all the same distance from the nucleus. They all have SIMILAR (but not necessarily the SAME) energy.

- Shells are numbered (1-... - Elements on the periodic table have shells numbered from 1 to 7)
- Higher numbers correspond to greater distance from the nucleus and greater energy, and larger size!
- Higher shells can hold more electrons than lower shells!




SUBSHELLS: Within a SHELL, electrons may move in different ways around the nucleus! These different "paths" are called SUBSHELLS


- SHAPES of regions of space that electrons are able to exist in.



"s" subshell  
(a spherical region)



"p" subshell  
(a dumbbell shaped region)



"d" subshell

- Some atoms also have "f" subshells (not pictured)

See p 248-251 for nicer drawings of the subshells.

ORBITALS - are specific regions of space where electrons may exist

- The SHAPE of an orbital is defined by the SUBSHELL it is in
- The ENERGY of an orbital is defined by both the SHELL the orbital is in AND the kind of SUBSHELL it is in
- Each orbital may, at most, contain TWO ELECTRONS

## ARRANGEMENT OF SHELLS, SUBSHELLS, AND ORBITALS

- Shells are numbered. Each shell can contain the same number of SUBSHELLS as its number:

1st shell: ONE possible subshell (s)


2nd shell: TWO possible subshells (s, p)

3rd shell: THREE possible subshells (s, p, d)

4th shell: FOUR possible subshells (s, p, d, f)




... and so on

- Each subshell can contain one or more ORBITALS, depending on how many different ways there are to arrange an orbital of that shape around the nucleus.



"s" subshell  
One possible  
orientation

Maximum 2 electrons in 1 orbital



"p" subshell: Three possible orientations

Maximum 6 electrons in 3 orbitals

- There are five possible orbitals in a "d" subshell, and 7 possible orbitals in an "f" subshell!

Maximum 10 electrons  
in 5 orbitals  
(see p 251)

Maximum 14 electrons  
in 7 orbitals

## ENERGY DIAGRAM

- We can map out electrons around an atom using an energy diagram:

