


CHEMICAL CALCULATIONS CONTINUED: REACTIONS

- Chemical reactions proceed on an ATOMIC basis, NOT a mass basis!
- To calculate with chemical reactions (i.e. use chemical equations), we need everything in terms of ATOMS ... which means MOLES of atoms




coefficients are in terms of atoms and molecules!

$$2 \text{ atoms Al} = 3 \text{ molecules Br}_2 = 2 \text{ formula units AlBr}_3$$

$$2 \text{ mol Al} = 3 \text{ mol Br}_2 = 2 \text{ mol AlBr}_3$$

- To do chemical calculations, we need to:
 - Relate the amount of substance we know (mass or volume) to a number of moles
 - Relate the moles of one substance to the moles of another using the equation
 - Convert the moles of the new substance to mass or volume as desired



* Given that we have 25.0 g of liquid bromine, how many grams of aluminum would we need to react away all of the bromine?

① Convert grams of bromine to moles: Need formula weight $\text{Br}_2 : \frac{2 \times 79.90}{159.80}$

$$159.80 \text{ g Br}_2 = \text{mol Br}_2$$

$$25.0 \text{ g Br}_2 \times \frac{\text{mol Br}_2}{159.80 \text{ g Br}_2} = 0.15645 \text{ mol Br}_2$$

② Use the chemical equation to relate moles of bromine to moles of aluminum

$$2 \text{ mol Al} = 3 \text{ mol Br}_2$$

$$0.15645 \text{ mol Br}_2 \times \frac{2 \text{ mol Al}}{3 \text{ mol Br}_2} = 0.10430 \text{ mol Al}$$


③ Convert moles aluminum to mass: Need formula weight $\text{Al} : 26.98$

$$26.98 \text{ g Al} = \text{mol Al}$$

$$0.10430 \text{ mol Al} \times \frac{26.98 \text{ g Al}}{\text{mol Al}} = \boxed{2.81 \text{ g Al}}$$

Example:

How many milliliters of 6.00M hydrochloric acid (FW=36.46 g/mol) is needed to completely react with 25.0 g of sodium carbonate (FW=105.99 g/mol)?



1 - Convert 25.0 g sodium carbonate to moles. Use FORMULA WEIGHT.

2 - Convert moles sodium carbonate to moles HCl. Use CHEMICAL EQUATION

3 - Convert moles HCl to volume HCl solution. Use MOLARITY.

$$\textcircled{1} \quad 105.99 \text{ g Na}_2\text{CO}_3 = 1 \text{ mol Na}_2\text{CO}_3$$


$$25.0 \text{ g Na}_2\text{CO}_3 \times \frac{1 \text{ mol Na}_2\text{CO}_3}{105.99 \text{ g Na}_2\text{CO}_3} = 0.2358713086 \text{ mol Na}_2\text{CO}_3$$

$$\textcircled{2} \quad 2 \text{ mol HCl} = 1 \text{ mol Na}_2\text{CO}_3$$

$$0.2358713086 \text{ mol Na}_2\text{CO}_3 \times \frac{2 \text{ mol HCl}}{1 \text{ mol Na}_2\text{CO}_3} = 0.4717426172 \text{ mol HCl}$$

102 Example:

How many milliliters of 6.00M hydrochloric acid (FW=36.46 g/mol) is needed to completely react with 25.0 g of sodium carbonate (FW=105.99 g/mol)?



1 - Convert 25.0 g sodium carbonate to moles. Use FORMULA WEIGHT.

2 - Convert moles sodium carbonate to moles HCl. Use CHEMICAL EQUATION

3 - Convert moles HCl to volume HCl solution. Use MOLARITY.


$$\textcircled{3} \quad 6.00 \text{ mol HCl} = \text{L}$$

$$0.4717426172 \text{ mol HCl} \times \frac{\text{L}}{6.00 \text{ mol HCl}} = 0.0786 \text{ L of 6.00M HCl}$$

Since the problem asks for volume in mL, we need to convert the units of the answer:

$$\text{mL} = 10^{-3} \text{ L}$$

$$0.0786 \text{ L} \times \frac{\text{mL}}{10^{-3} \text{ L}} = \boxed{78.6 \text{ mL of 6.00M HCl}}$$



Calculate how many grams of acrylonitrile could be obtained from 651 g of propylene, assuming there is excess NO present.

- 1 - Convert 651 g propylene to moles. Use FORMULA WEIGHT.
- 2 - Convert moles propylene to moles acrylonitrile. Use CHEMICAL EQUATION
- 3 - Convert moles acrylonitrile to mass. Use FORMULA WEIGHT.


$$\textcircled{1} \quad 42.081 \text{ g C}_3\text{H}_6 = \text{mol C}_3\text{H}_6$$

$$\textcircled{2} \quad 4 \text{ mol C}_3\text{H}_6 = 4 \text{ mol C}_3\text{H}_3\text{N}$$

$$\textcircled{3} \quad 53.064 \text{ g C}_3\text{H}_3\text{N} = \text{mol C}_3\text{H}_3\text{N}$$

$$651 \text{ g C}_3\text{H}_6 \times \overset{\textcircled{1}}{\frac{\text{mol C}_3\text{H}_6}{42.081 \text{ g C}_3\text{H}_6}} \times \overset{\textcircled{2}}{\frac{4 \text{ mol C}_3\text{H}_3\text{N}}{4 \text{ mol C}_3\text{H}_6}} \times \overset{\textcircled{3}}{\frac{53.064 \text{ g C}_3\text{H}_3\text{N}}{\text{mol C}_3\text{H}_3\text{N}}} = 821 \text{ g C}_3\text{H}_3\text{N}$$

151.90 g/mol



How many mL of 0.250M potassium permanganate are needed to react with 3.36 g of iron(II) sulfate?

- 1 - Convert 3.36 g iron(II) sulfate to moles. Use FORMULA WEIGHT.
- 2 - Convert moles iron(II) sulfate to moles potassium permanganate. Use CHEMICAL EQUATION.
- 3 - Convert moles potassium permanganate to volume solution. Use MOLARITY.

$$\textcircled{1} 151.90 \text{ g FeSO}_4 = \text{mol FeSO}_4$$

$$\textcircled{2} 10 \text{ mol FeSO}_4 = 2 \text{ mol KMnO}_4$$

$$\textcircled{3} 0.250 \text{ mol KMnO}_4 = \text{L}$$

$$3.36 \text{ g FeSO}_4 \times \frac{\textcircled{1} \text{ mol FeSO}_4}{151.90 \text{ g FeSO}_4} \times \frac{\textcircled{2} 2 \text{ mol KMnO}_4}{10 \text{ mol FeSO}_4} \times \frac{\textcircled{3} \text{ L}}{0.250 \text{ mol KMnO}_4} = 0.0177 \text{ L}$$


Convert the final answer to mL as directed in the problem statement,

$$\text{mL} = 10^{-3} \text{ L}$$

$$0.0177 \text{ L} \times \frac{\text{mL}}{10^{-3} \text{ L}} = \boxed{17.7 \text{ mL of } 0.250 \text{ M KMnO}_4}$$

CONCEPT OF LIMITING REACTANT

- When does a chemical reaction STOP?



- When does this reaction stop? When burned in open air, this reaction stops when all the MAGNESIUM STRIP is gone. We say that the magnesium is LIMITING.

- This reaction is controlled by the amount of available magnesium

- At the end of a chemical reaction, the LIMITING REACTANT will be completely consumed but there may be amount of OTHER reactants remaining. We do chemical calculations in part to minimize these "leftovers".

These are often called "excess" reactants, or reactants present "in excess"