$$\frac{(2x)^2}{(0.10625-x)(0.10625-x)} = 0.0123$$

You can solve this problem with the quadratic

$$\frac{2x}{0.10625^{-x}} = 0.1109053651$$

$$2x = (0.1109053651)(0.16625-x)$$

$$18.03339269x = 0.10625-x$$

$$19.03339269x = 0.10625$$

$$x = 0.0055822943$$

Species	[cavil.]
N2	0.10625 - x
01	0.10625-x
NO	2x

Now, use the value of 'x' to find the concentrations:


$$N_2$$
: 0.10625 -  $x = 0.101M$   
 $O_2$ : 0.10625 -  $y = 0.101M$   
 $N_0$ :  $2x = 0.0112M$ 

We know Kc has a small value (0.0123), which suggests that concentration of NO (product) should be small relative to reactant concentration.


- Pressure can affect a GAS-PHASE equilibrium ... sometimes. How?

$$(O(g) + 2H_2(g) \rightleftharpoons CH_3OH(g)$$

- ... how might pressure affect this equilibrium?
- If the change in pressure CHANGES CONCENTRATIONS, then this equilibrium would be disturbed and Le Chateleir's Principle would apply.
  - Adding an INERT GAS would change pressure, but would it change concentration of the gases? NO so addition of argon would have no effect on the equilibrium!
    - What about COMPRESSION?



... compression increases pressure by DECREASING total volume.



... but this volume change affects ALL concentrations the same way. In this example, each concentration is DOUBLED.

$$(O(g) + 2H_2(g) \rightleftharpoons (H_3OH(g))$$

$$(I) = \frac{(I)}{(I)(I)^2} = \frac{(I)}{(I)(I)^2}$$
For simplicity, let's assume Kc = 1, and all concs = 1M

$$\frac{Doubling}{gives Q = (2)(2)^2} = \frac{1}{4}$$

Q < Kc, so equilibrium shifts to the RIGHT, forming more methanol at the expense of hydrogen and carbon monoxide.

In general, compressing an equilibrium reaction in the gas phase will cause the equilibrium to shift towards the side with fewer moles of gas. This causes the pressure to decrease.

In general, decompressing an equilibrium reaction in the gas phase will cause the equilibrium to shift towards the side with more moles of gas. This causes the pressure to increase.

HOWEVER, this can only be true IF there's a side of the reaction with more moles of gas than the other. If both sides of the reaction have the SAME number of moles of gas, then a pressure change will NOT affect the equilibrium.

Example: 
$$N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$$

... would not respond to a pressure change.

# 119 FACTORS THAT MAY AFFECT EQUILBRIUM

1) TEMPERATURE (effect depends on whether reaction is endothermic or exothermic)


- Changes rate of reaction, too!

... changes Kc

PRESSURE - only for gas-phase reactions which have different numbers of moles of gas on each side of the equilbrium. Otherwise, no effect.

... no change of Kc

(3) CATALYSTS - do NOT affect equilibrium, but make the equilbrium state occur more quickly.



The catalyst raises BOTH forward and reverse rates, so it doesn't affect the composition of the equilibrium mixture!



CONCENTRATION - Le Chateleir's Principle applies for changing concentrations. An equilibrium will shift to counteract a change in concentration of reactant or product.

... doesn't change Kc.

## ACID/BASE EQUILIBRIUM

- Several scientific theories exist that define acid-base chemistry. We will discuss THREE of these theories.
- These theories differ in the way that acids, bases, and their associated reactions are defined.
- Typically, the newer theories include MORE chemicals under the umbrella of "acid-base chemistry"!

#### THREE ACID-BASE THEORIES

- (I) Arrhenius theory
- (2) Bronsted-Lowry theory
- (3) Lewis theory

- The oldest model of acid-base chemistry!

- Only applicable to systems where WATER is the solvent!

ACIDS are substances that ionize in water to increase the concentration of HYDRONIUM ION

$$HA + H_2O = H_3O^{\dagger} + A^{-}$$
Hydronium ion

or, for simplicity: 
$$HA \stackrel{H_2O}{\longrightarrow} H^{\dagger} + A^{-}$$

"Hydrogen ion" - doesn't really exist as a free ion in water, but a convenient simplification!

# BASES are substances that ionize in water to increase the concentration of HYDROXIDE ION

For soluble metal hydroxides:  $V_{\alpha}O_{H} \rightarrow V_{\alpha}^{+} + O_{H}^{-}$   $MOH \stackrel{HzO}{\longleftarrow} M^{+} + OH^{-}$ Hydroxide ion


For other Arrhenius bases:

An Arrhenius acid base reaction can be represented by:

$$H_3O^+ + OH^- \rightleftharpoons 2H_2O$$
 "neutralization"

or, using hydrogen ion instead of hydronium


#### **BRONSTED-LOWRY THEORY**




- Bronsted-Lowry theory views acid-base reactions as <u>PROTON</u> TRANSFER reactions!

#### **ACIDS are PROTON DONORS**


#### **BASES are PROTON ACCEPTORS**



A CONJUGATE PAIR is an acid and a base that differ by a proton!



... you should be able to write the products of a Bronsted-Lowry acid-base reaction, identifying the CONJUGATE PAIRS



B+H20 
$$=$$
 BH++OH-
base conjugate acid
of B

NH3+IH20  $=$  NH4+OH-
This is why we often call an ammonia/water solution "ammonium hydroxide"!

In the red reactions, water functions as a base. In the blue reactions, water functions as a acid!

### LEWIS THEORY


- Lewis theory treats acid-base chemistry as ELECTRON-TRANSFER chemistry involving pairs of electrons
- Lewis acid-base reactions form new covalent bonds (of interest to organic chemists!)

## ACIDS are ACCEPTORS of electron pairs

... this is why some METAL IONS, even though they contain no hydorgen ions, can exhibit ACIDIC character. Many metal ions can accept a pair of electrons to form a COMPLEX with a Lewis base! ex,  $h_a(NH_3)_3^+$ 

BASES are DONORS of electron pairs.

... so, Lewis bases have LONE PAIRS OF ELECTRONS in their Lewis structures



... In a Lewis acid-base reaction, electrons are donated from the Lewis base to the Lewis acid. This forms a new COVALENT BOND between the acid and the base.

## Example: ammonia and water


Overall: 
$$H_2O + NH_3 \rightleftharpoons NH_4^{\dagger} + OH^-$$

- From Arrhenius to Lewis, the definitions get broader as you go along. In other woeds, the later definitions include MORE SUBSTANCES under the acid/base umbrella.

If something is an Arrhenius acid, it is also an acid in the Bronsted or Lewis picture. If something is an Arrhenius base, it is also a base in the Bronsted or Lewis picture.

All Bronsted acids are Lewis acids, and all Bronsted bases are Lewis bases.

... but not all Lewis acids/bases (like the metal ions) are Bronsted or Arrhenius acids/bases.



... We will primarily use the <u>BRONSTED-LOWRY</u> theory from this point in the course!

- Water self-ionizes!

$$2 H_{2}O \rightleftharpoons H_{3}O^{+} OH^{-}$$
or
$$H_{2}O \rightleftharpoons H^{+} + OH^{-}$$

This is an equilibrium reaction!

$$K = \frac{[H_3D^+][OH^-]}{[H_2O]^2}$$
(X) = molar concentration of "X"

In aqueous solution, ( $H_2D$ ) is essentially constant, so we roll that into K.

$$K_{w} = [H_{3}D^{+}][OH^{-}] = 1.0 \times 10^{-14}$$

This is the value at 25C