

¹⁵⁸ SINCE the enthalpy change does NOT depend on path, this means that we can use standard values for enthalpy to predict the heat change in reactions that we have not tested in a calorimeter.

THERMOCHEMICAL EQUATIONS

- is like a regular chemical equation, except that phase labels are REQUIRED and the enthalpy for the reaction is given along with the equation.



- Why are phase labels required? Because phase changes either absorb or release energy.

$\Delta H = -1800 \text{ kJ}$... what does this mean?

$$1 \text{ mol CH}_3\text{COCH}_3 = -1800 \text{ kJ}$$

$$4 \text{ mol O}_2 = -1800 \text{ kJ}$$

$$3 \text{ mol CO}_2 = -1800 \text{ kJ}$$

$$3 \text{ mol H}_2\text{O} = -1800 \text{ kJ}$$

We treat the enthalpy change as if it's another product of the reaction!



What would be the enthalpy change when 25 g of water are produced by the reaction?

1 - Convert 25 g water to moles. Use FORMULA WEIGHT.

2 - Convert moles water to enthalpy change. Use THERMOCHEMICAL EQUATION

$$\textcircled{1} \text{H}_2\text{O} - \text{H} : 2 \times 1.008$$

$$\text{O} : 1 \times 16.00$$

$$\hline 18.016 \text{ g H}_2\text{O} = \text{mol H}_2\text{O}$$

$$\textcircled{2} 3 \text{ mol H}_2\text{O} = -1800 \text{ kJ}$$

$$25 \text{ g H}_2\text{O} \times \frac{\textcircled{1} \text{ mol H}_2\text{O}}{18.016 \text{ g H}_2\text{O}} \times \frac{\textcircled{2} -1800 \text{ kJ}}{3 \text{ mol H}_2\text{O}} = \boxed{-830 \text{ kJ}}$$

Notes: 1) This reaction is EXOTHERMIC ("- sign for enthalpy/heat). Energy is released!


2) If this is a constant pressure process, Q also = -830 kJ

A few more terms related to enthalpy:

- Enthalpy of vaporization / heat of vaporization: The enthalpy change on vaporizing one mole of a substance. (from liquid to vapor)

- Enthalpy of fusion / heat of fusion: The enthalpy change when a mole of liquid changes to the solid state.

 Phase changes require energy, too!




Calculate the enthalpy change for the combustion of 1.00 kg of hydrogen gas.

1 - Convert 1.00 kg hydrogen gas to moles. Use FORMULA WEIGHT.

2 - Convert moles hydrogen gas to enthalpy change. Use THERMOCHEMICAL EQUATION

$$\textcircled{1} \quad 2.016 \text{ g H}_2 = 1 \text{ mol H}_2 \quad \text{kg} = 10^3 \text{ g} \quad \textcircled{2} \quad 2 \text{ mol H}_2 = -484 \text{ kJ}$$

$$1.00 \text{ kg H}_2 \times \frac{10^3 \text{ g}}{\text{kg}} \times \frac{\overset{\textcircled{1}}{1 \text{ mol H}_2}}{2.016 \text{ g H}_2} \times \frac{\overset{\textcircled{2}}{-484 \text{ kJ}}}{2 \text{ mol H}_2} = \boxed{-120000 \text{ kJ per kg H}_2}$$



What is the enthalpy change when 150. L of nitrogen monoxide are formed by this reaction at 25.0 C and 1.50 atm pressure?

1 - Convert 150. L NO to moles. Use IDEAL GAS EQUATION.

2 - Convert moles NO to enthalpy change. Use THERMOCHEMICAL EQUATION.

$$\textcircled{1} PV = nRT \quad \left| \quad P = 1.50 \text{ atm} \quad V = 150. \text{ L}$$


$$n = \frac{PV}{RT} \quad \left| \quad R = 0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}} \quad T = 25.0^\circ\text{C} = 298.2 \text{ K}$$

$$n_{\text{NO}} = \frac{(1.50 \text{ atm})(150. \text{ L})}{\left(0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}}\right)(298.2 \text{ K})} = 9,194,822,849 \text{ mol NO}$$

$$\textcircled{2} 4 \text{ mol NO} = -906 \text{ kJ}$$

$$9,194,822,849 \text{ mol NO} \times \frac{-906 \text{ kJ}}{4 \text{ mol NO}} = \boxed{-2080 \text{ kJ}}$$

propane



Calculate the volume of propane gas at 25.0 C and 1.08 atm required to provide 565 kJ of heat using the reaction above.

- 1 - Convert 565 kJ to moles propane. Use THERMOCHEMICAL EQUATION
- 2 - Convert moles propane to volume. Use IDEAL GAS EQUATION.

$$\textcircled{1} \text{ mol C}_3\text{H}_8 = -2043 \text{ kJ}$$

Note: The 565 kJ of energy is supplied by the reaction. That means the reaction loses 565 kJ, and since the calculation is from the point of view of the reaction, we start with -565 kJ.

$$-565 \text{ kJ} \times \frac{\text{mol C}_3\text{H}_8}{-2043 \text{ kJ}} = 0.2765540871 \text{ mol C}_3\text{H}_8$$

$$\textcircled{2} \text{ PV} = nRT \quad \left| \quad n = 0.2765540871 \text{ mol C}_3\text{H}_8 \quad R = 0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}} \right.$$

$$V = \frac{nRT}{P} \quad \left| \quad T = 25.0^\circ\text{C} = 298.2 \text{ K} \quad P = 1.08 \text{ atm} \right.$$

$$V = \frac{(0.2765540871 \text{ mol C}_3\text{H}_8) (0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}}) (298.2 \text{ K})}{(1.08 \text{ atm})}$$

$$= 6.27 \text{ L C}_3\text{H}_8 @ 25.0^\circ\text{C}, 1.08 \text{ atm}$$

END OF CHAPTER 6