


Example:

How many milliliters of 6.00M hydrochloric acid is needed to completely react with 25.0 g of sodium carbonate?



1 - Convert 25.0 g sodium carbonate to moles. Use FORMULA WEIGHT.

2 - Convert moles sodium carbonate to moles HCl. Use CHEMICAL EQUATION

3 - Convert moles HCl to volume solution. Use MOLARITY

$$\textcircled{1} \text{Na}_2\text{CO}_3 - \text{Na} : 2 \times 22.99$$

$$\text{C} : 1 \times 12.01$$

$$\text{O} : 3 \times 16.00$$


$$\underline{\underline{105.99 \text{ g Na}_2\text{CO}_3 = \text{mol Na}_2\text{CO}_3}}$$

$$25.0 \text{ g Na}_2\text{CO}_3 \times \frac{\text{mol Na}_2\text{CO}_3}{105.99 \text{ g Na}_2\text{CO}_3} = 0.2358713086 \text{ mol Na}_2\text{CO}_3$$

$$\textcircled{2} 2 \text{ mol HCl} = \text{mol Na}_2\text{CO}_3$$

$$0.2358713086 \text{ mol Na}_2\text{CO}_3 \times \frac{2 \text{ mol HCl}}{\text{mol Na}_2\text{CO}_3} = 0.4717426172 \text{ mol HCl}$$

How many milliliters of 6.00M hydrochloric acid is needed to completely react with 25.0 g of sodium carbonate?



-
- 1 - Convert 25.0 g sodium carbonate to moles. Use FORMULA WEIGHT.
 - 2 - Convert moles sodium carbonate to moles HCl. Use CHEMICAL EQUATION
 - 3 - Convert moles HCl to volume solution. Use MOLARITY
-


$$\textcircled{3} \quad 6.00 \text{ mol HCl} = \text{L}$$

$$0.4717426172 \text{ mol HCl} \times \frac{\text{L}}{6.00 \text{ mol HCl}} = 0.0786 \text{ L}$$

... but since the problem statement wants the answer in mL ...

$$\text{mL} = 10^{-3} \text{ L}$$

$$0.0786 \text{ L} \times \frac{\text{mL}}{10^{-3} \text{ L}} = \boxed{78.6 \text{ mL of } 6.00 \text{ M HCl}}$$



Calculate how many grams of acrylonitrile could be obtained from 651 g of propylene, assuming there is excess NO present.

- 1 - Convert 651 g propylene to moles using FORMULA WEIGHT.
- 2 - Convert moles propylene to moles acrylonitrile. Use CHEMICAL EQUATION
- 3 - Convert moles acrylonitrile to grams acrylonitrile. Use FORMULA WEIGHT.


$$\textcircled{1} \quad 42.081 \text{ g C}_3\text{H}_6 = \text{mol C}_3\text{H}_6$$

$$\textcircled{2} \quad 4 \text{ mol C}_3\text{H}_6 = 4 \text{ mol C}_3\text{H}_3\text{N}$$

$$\textcircled{3} \quad 53.064 \text{ g C}_3\text{H}_3\text{N} = \text{mol C}_3\text{H}_3\text{N}$$

$$\begin{aligned}
 & 651 \text{ g C}_3\text{H}_6 \times \frac{\text{mol C}_3\text{H}_6}{42.081 \text{ g C}_3\text{H}_6} \times \frac{4 \text{ mol C}_3\text{H}_3\text{N}}{4 \text{ mol C}_3\text{H}_6} \times \frac{53.064 \text{ g C}_3\text{H}_3\text{N}}{\text{mol C}_3\text{H}_3\text{N}} \\
 & = \boxed{821 \text{ g C}_3\text{H}_3\text{N}}
 \end{aligned}$$

151.90 g/mol



How many mL of 0.250M potassium permanganate are needed to react with 3.36 g of iron(II) sulfate?

- 1 - Convert 3.36 g iron(II) sulfate to moles. Use FORMULA WEIGHT.
- 2 - Convert moles iron(II) sulfate to moles potassium permanganate. Use CHEMICAL EQUATION.
- 3 - Convert moles potassium permanganate to volume of solution. Use MOLARITY.

$$\textcircled{1} 151.90 \text{ g FeSO}_4 = \text{mol FeSO}_4$$

$$\textcircled{2} 10 \text{ mol FeSO}_4 = 2 \text{ mol KMnO}_4$$

$$\textcircled{3} 0.250 \text{ mol KMnO}_4 = \text{L}$$

$$3.36 \text{ g FeSO}_4 \times \frac{\text{mol FeSO}_4}{151.90 \text{ g FeSO}_4} \times \frac{2 \text{ mol KMnO}_4}{10 \text{ mol FeSO}_4} \times \frac{\text{L}}{0.250 \text{ mol KMnO}_4} = 0.0177 \text{ L of } 0.250 \text{ M KMnO}_4$$


As in a previous example, we need the final volume in mL ...

$$\text{mL} = 10^{-3} \text{ L}$$

$$0.0177 \text{ L} \times \frac{\text{mL}}{10^{-3} \text{ L}} = \boxed{17.7 \text{ mL of } 0.250 \text{ M KMnO}_4}$$

CONCEPT OF LIMITING REACTANT

- When does a chemical reaction STOP?




- When does this reaction stop? When burned in open air, this reaction stops when all the MAGNESIUM STRIP is gone. We say that the magnesium is LIMITING.
- This reaction is controlled by the amount of available magnesium
- At the end of a chemical reaction, the LIMITING REACTANT will be completely consumed, but there may be amount of OTHER reactants remaining. We do chemical calculations in part to minimize these "leftovers".

These are often called "excess" reactants, or reactants present "in excess"

LIMITING REACTANT CALCULATIONS

- To find the limiting reactant, calculate how much product would be produced from ALL given reactants. Whichever produces the SMALLEST amount of product is the limiting reactant, and the smallest amount of product is the actual amount of product produced.



If you start with 100. g of each reactant, how much calcium carbide would be produced?

① $56.08 \text{ g CaO} = \text{mol CaO}$ ② $\text{mol CaO} = \text{mol CaC}_2$ ③ $64.10 \text{ g CaC}_2 = \text{mol CaC}_2$

$$100. \text{g CaO} \times \frac{\text{mol CaO}}{56.08 \text{g CaO}} \times \frac{\text{mol CaC}_2}{\text{mol CaO}} \times \frac{64.10 \text{g CaC}_2}{\text{mol CaC}_2} = 114 \text{ g CaC}_2$$

① $12.01 \text{ g C} = \text{mol C}$ ② $3 \text{ mol C} = \text{mol CaC}_2$ ③ $64.10 \text{ g CaC}_2 = \text{mol CaC}_2$

$$100. \text{g C} \times \frac{\text{mol C}}{12.01 \text{g C}} \times \frac{\text{mol CaC}_2}{3 \text{ mol C}} \times \frac{64.10 \text{g CaC}_2}{\text{mol CaC}_2} = 178 \text{ g CaC}_2$$

This reaction stops when it reaches 114 g of calcium carbide produced. At that point, there is no longer any CaO for the carbon to react with. We say that CaO is LIMITING, and carbon is present IN EXCESS.

PERCENT YIELD

- Chemical reactions do not always go to completion! Things may happen that prevent the conversion of reactants to the desired/expected product!


① SIDE REACTIONS:



... so in a low-oxygen environment, you may produce less carbon dioxide than expected!

② TRANSFER AND OTHER LOSSES

- When isolating a product, losses may occur in the process. Example: filtering



③ EQUILIBRIUM

- Reactions may reach an equilibrium between products and reactants. We'll talk more about this in CHM 111. The net result is that the reaction will appear to stop before all reactants have been consumed!

- All of these factors cause a chemical reaction to produce LESS product than calculated. For many reactions, this difference isn't significant. But for others, we need to report the PERCENT YIELD.

$$\text{PERCENT YIELD} = \frac{\text{ACTUAL YIELD}}{\text{THEORETICAL YIELD}} \times 100\%$$

↙ Determined EXPERIMENTALLY!

↑ Calculated based on the limiting reactant. (The chemical calculations you've done up to now have been theoretical yields!)

... the percent yield of a reaction can never be greater than 100% due to conservation of mass! If you determine that a percent yield is greater than 100%, then you've made a mistake somewhere - either in a calculation or in the experiment itself!