

- a SOLUTION is a HOMOGENEOUS MIXTURE.

- parts of a solution:

- component(s) of a solution present in small amounts.

2)SOLVENT

- the component of a solution present in the GREATEST amount

- in solutions involving a solid or gas mixed with a LIQUID, the liquid is typically considered the solvent.

- solutions are usually the same phase as the pure solvent. For example, at room temperature salt water is a liquid similar to pure water.

#### 55 SOLVENTS

- We traditionally think of solutions as involving gases or solids dissolved in liquid solvents. But ANY of the three phases may act as a solvent!

() GAS SOLVENTS

- Gases are MISCIBLE, meaning that they will mix together in any proportion.
- This makes sense, since under moderate conditions the molecules of a gas don't interact wth each other.
- Gas solvents will only dissolve other gases.

2) LIQUID SOLVENTS

- Can dissolve solutes that are in any phase: gas, liquid, or solid.
- Whether a potential solute will dissolve in a liquid depends on how compatible the forces are between the liquid solvent and the solute.

## 3 SOLID SOLVENTS

- Solids can dissolve other solids, and occasionally liquids.
- Solid-solid solutions are called ALLOYS. Brass (15% zinc dissolved in copper) is a good example.
- AMALGAM is a solution resulting from dissolving mercury into another metal.


#### HOW THINGS DISSOLVE

- Let's look at how things dissolve into water, since aqueous solutions are quite common.

sucrose (table sugar)  

$$C_{12}H_{22}O_{11}(s) \xrightarrow{H_20} (_{12}H_{22}O_{11}(a_q))$$

... what happens?



- Water molecules pull the sugar molecules out of the sugar crystal and into solution.

- Attractions between sugar molecules and water allow this to happen.


- The solubility of the sugar depends on how well water and sugar interact (HYDRATION) versus how well the sugar molecules are held in the crystal (LATTICE ENERGY)

- "like dissolves like": Substances held together by similar (or at least compatible) kinds of attractive forces can dissolve in each other. Substances that are held together by very different kinds of attractive forces will not dissolve in one another!

Consider WATER:


HYDROGEN BONDS

Water mixes well with other substances that can hydrogen bond, like ETHANOL!



Water can dissolve polar substances! (SUCROSE is polar!)

н н н н - С-С-О-Н-О-Н н н



Hydrogen bond between ethanol and water


Since IONIC BONDS are also interactions between opposite charges (You can think of an ionic bond here as an extreme case of dipole-dipole interaction), many IONIC SUBSTANCES will

also dissolve in water!

SMALL (little London force)  $\sqrt{}$ 

large and/or nonpolar solutes do not dissolve well in water!

(example: OILS and WAXES)



#### MOLECULAR AND IONIC SOLUTIONS

- MOLECULAR solutions:

Contain MOLECULES dissolved in one another.

1 - Any mixture of GASES

- all gases mix with one another, since gas molecules (effectively) do not interact with one another.

<u> </u> - Liquids

- Liquids dissolve well in one another only if they are held together by similar kinds of forces

### 3 - Solids and liquids


- MOLECULAR SOLIDS will dissolve well in liquids if they are held together by similar forces.

- IONIC SOLIDS will sometimes dissolve in POLAR liquids, but not in nonpolar liquids

- COVALENT NETWORK solids don't generally dissolve well in other substances

## IONIC solutions

- form when ions from IONIC SUBSTANCES interact with POLAR solvents - often WATER.



The charged ends of the water molecule HYDRATE the ions.

- The solubility of an ionic compound depends on whether HYDRATION (attraction of water molecules for an ion) is greater than LATTICE ENERGY - the attraction of ions in a crystal lattice for one another..

- SMALLER IONS are usually easier to enclose in water than larger ones, and ions with larger charges are attracted to water molecules.

- But solubility is also determined by LATTICE ENERGY - which holds the solid ionic compound together. Ions with high charges tend to be strongly attracted to other ions in a crystal, meaning lattice energy is high. Smaller ions also tend to have higher lattice energies. Lattice energy and hydration are competing trends!

#### EXTERNAL FACTORS AFFECTING SOLUBILITY

- There are a few external factors that affect the solubility. (By external, we mean other than the chemical identity of the solute and solvent).

## 

-For gases dissolved in liquids, the solubility DECREASES as the temperature INCREASES

- This is why THERMAL POLLUTION is bad! Hot water holds less oxygen than cooler water.

- For solids dissolved in liquids, solubility USUALLY increases with temperature. This is not true for ALL solid/liquid solutions.

# D PRESSURE

- For gases dissolved in liquids, solubility INCREASES when the partial pressure of the solute gas over the solution INCREASES.

- Consider soft drinks. They go flat after opening because the pressure of carbon dioxide over the liquid goes down.

- No significant pressure effects for solid/liquid solutions.

# CONCENTRATION

61

- When you discuss a solution, you need to be aware of:

- what materials are in the solution

- how much of each material is in the solution

- CONCENTRATION is the amount of one substance compared to the others in a solution. This sounds vague, but that's because there are many different ways to specify concentration!

- We will discuss four different concentration units in CHM 111:

How would you prepare 455 grams of an aqueous solution that is 6.50% sodium sulfate by mass?

$$\frac{mass \%}{16.SD} = \frac{mass Na_2 Soy}{mass Solution} \times 100$$

Start any concentration calculation by writing out the DEFINITION of each concentration unit you're using...

We know everytihng in the definition EXCEPT mass sodium sulfate. So calculate the mass of sodium sulfate required to prepare the solution.

$$6.50 = \frac{muss N_{2}SOy}{4SSg} \times 100$$

$$\frac{10 \times 43Sg}{29.6g} = \frac{100}{29.6g} \times 100$$

We also need the amount of water. Find it by subtraction ...

So mix 29.6 grams sodium sulfate and 425.4 grams water to prepare the solution.

What's the MOLALITY and MOLE FRACTION OF SOLUTE of the previous solution?

29.6 g 
$$N_{42}SO_{4}$$
, 425.4 g water  $\leftarrow$  previous solution  
 $m = \frac{mol Na_2SO_{4}}{Kg H_2O}$  (solute) (1)  
(solvent) (2)

1 - Convert 29.6 grams sodium sulfate to moles. Use FORMULA WEIGHT.

2 - Convert 425.4 grams water to kg.

$$Na_{2}So_{4} = Na - 2x22.99$$
  
 $S - 1 x32.07$   
 $O - 4x 16.00$   
 $142.0Sg Na_{2}So_{4} = mol Na_{2}So_{4}$ 

$$29.6 g N_{62} S_{04} \times \frac{m_0 N_{62} S_{04}}{142.05 g N_{62} S_{04}} = 0.20837733 | 9 m_0 N_{62} S_{04}$$

$$K_g = 10^3 g \qquad (2)$$

$$H_2 S_{04} \times \frac{K_g}{10^3 g} = 0.4254 K_g H_{20}$$

$$= 0.4254 K_g H_{20}$$

$$= 0.490 m N_{62} S_{04}$$