

2500 L of chlorine gas at 25.0 C and 1.00 atm are used to make hydrochloric acid. How many kilograms of hydrochloric acid could be produced if all the chlorine reacts?



- 1 - Convert 2500 L chlorine gas to moles. Use IDEAL GAS EQUATION.
- 2 - Convert moles chlorine gas to moles HCl. Use CHEMICAL EQUATION
- 3 - Convert moles HCl to mass HCl. Use FORMULA WEIGHT. (and a g \rightarrow kg conversion)

$$\textcircled{1} \begin{array}{l} PV = nRT \\ n = \frac{PV}{RT} \end{array} \quad \begin{array}{l} P = 1.00 \text{ atm} \quad R = 0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}} \\ V = 2500 \text{ L} \quad T = 25.0^\circ\text{C} = 298.2 \text{ K} \end{array}$$

$$n_{\text{Cl}_2} = \frac{(1.00 \text{ atm})(2500 \text{ L})}{(0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}})(298.2 \text{ K})} = 102.1646983 \text{ mol Cl}_2$$

$$\textcircled{2} \text{ mol Cl}_2 = 2 \text{ mol HCl} \quad \textcircled{3} 36.458 \text{ g HCl} = \text{mol HCl}$$

$$102.1646983 \text{ mol Cl}_2 \times \frac{2 \text{ mol HCl}}{\text{mol Cl}_2} \times \frac{36.458 \text{ g HCl}}{\text{mol HCl}} = 7450 \text{ g HCl}$$

$$\begin{array}{l} \text{HCl: H} - 1 \times 1.008 \\ \quad \text{Cl} - 1 \times 35.45 \\ \hline 36.458 \end{array}$$

Convert to kilograms (problem specifies these units for answer) $\text{Kg} = 10^3 \text{g}$

$$7450 \text{ g HCl} \times \frac{\text{Kg}}{10^3 \text{g}} = \boxed{7.45 \text{ Kg HCl}}$$



If 48.90 mL of 0.250 M HCl solution reacts with sodium carbonate to produce 50.0 mL of carbon dioxide gas at 290.2 K, what is the pressure of the carbon dioxide gas?

- 1 - Convert 48.90 mL of HCl solution to moles. Use MOLARITY.
- 2 - Convert moles HCl to moles carbon dioxide. Use CHEMICAL EQUATION.
- 3 - Convert moles carbon dioxide to pressure. Use IDEAL GAS EQUATION.

$$\textcircled{1} 0.250 \text{ mol HCl} = \text{L} \quad \text{mL} = 10^{-3} \text{ L} \quad \textcircled{2} 2 \text{ mol HCl} = \text{mol CO}_2$$

$$48.90 \text{ mL} \times \frac{10^{-3} \text{ L}}{\text{mL}} \times \frac{0.250 \text{ mol HCl}}{\text{L}} \times \frac{\text{mol CO}_2}{2 \text{ mol HCl}} = 0.0061125 \text{ mol CO}_2$$

$$\textcircled{3} \begin{array}{l} PV = nRT \\ P = \frac{nRT}{V} \end{array} \left| \begin{array}{l} n = 0.0061125 \text{ mol CO}_2 \\ R = 0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}} \end{array} \right. \begin{array}{l} T = 290.2 \text{ K} \\ V = 50.0 \text{ mL} = 0.0500 \text{ L} \end{array}$$

$$P = \frac{(0.0061125 \text{ mol CO}_2) \left(0.08206 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}} \right) (290.2 \text{ K})}{(0.0500 \text{ L})} =$$

$$= \boxed{2.91 \text{ atm}}$$

- thermodynamics: the study of energy transfer

Conservation of energy: Energy may change form, but the overall amount of energy remains constant. "first law of thermodynamics"

- ... but what IS energy?

- energy is the ability to do "work"

↑
motion of matter


Kinds of energy?

- Kinetic energy: energy of matter in motion $E_K = \frac{1}{2} m v^2$

mass

velocity

- Potential energy: energy of matter that is being acted on by a field of force (like gravity)



When the ball falls, its potential energy is converted to kinetic!

- What sort of energy concerns chemists? Energy that is absorbed or released during chemical reactions.

- Energy can be stored in chemicals ... molecules and atoms.

INTERNAL ENERGY: "U"



related to the kinetic and potential energy of atoms, molecules, and their component parts.

- We measure energy transfer ... which is called HEAT. (HEAT is the flow of energy from an area of higher temperature to an area of lower temperature)

Q: heat

SYSTEM: the object or material under study

SURROUNDINGS: everything else

Type of process	Energy is ...	Sign of Q	Temp of SURROUNDINGS ...
ENDOTHERMIC	transferred from SURROUNDINGS to SYSTEM	+	decreases
EXOTHERMIC	transferred from SYSTEM to SURROUNDINGS	-	increases