

A DOT STRUCTURE FOR A MOLECULE WITH DELOCALIZED BONDS

$$O = 3 \times 6 = 18$$

See text, 9.7
p 356-358

① Count valence electrons

② Pick central atom and draw skeletal structure


- central atom is usually the one that needs to gain the most electrons!

- skeletal structure has all atoms connected to center with single bonds

③ Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.

④ Check octet rule - each atom should have a share in 8 electrons (H gets 2). if not, make double or triple bonds.

O_3 (OZONE)




Central oxygen has only six electrons



The structure we drew implies that one of the outer oxygen atoms is closer to the central oxygen atom than the other one.

Experimentally, though, we find the two oxygen atoms to be the SAME distance from the center.

In the ozone molecule, electrons are actually being shared between ALL THREE oxygen atoms at the same time. This is called a DELOCALIZED BOND.



The structures in the green box are called RESONANCE STRUCTURES. The "real" structure of ozone is an "average" of the two resonance structures. The "double bond" electrons in these structures are actually shared between all three oxygen atoms

A DOT STRUCTURE FOR A POLYATOMIC ION





$$\text{N}: 1 \times 5$$

$$\text{H}: 4 \times 1$$

 9 electrons

$$- 1 \text{ electron (+1 charge)}$$

 8 valence electrons



To indicate the charge of the ion, enclose the structure in large brackets, then put the charge at the upper right. ... much like we would write the charge of other ions like magnesium...



- ① Count valence electrons
- ② Pick central atom and draw skeletal structure
 - central atom is usually the one that needs to gain the most electrons!
 - skeletal structure has all atoms connected to center with single bonds
- ③ Distribute remaining valence electrons around structure, outer atoms first. Follow octet rule until you run out of electrons.
- ④ Check octet rule - each atom should have a share in 8 electrons (H gets 2). if not, make double or triple bonds.

EXPANDED VALENCE and other exceptions to the "octet rule"

- Some atoms do not always obey the octet rule. A few, like BORON, will bond in such a way that they end up with less than eight electrons.





... but many more bond in such a way that they end up with a share in MORE THAN EIGHT electrons!

- Any atom in period three or greater can do this. SULFUR and PHOSPHORUS compounds commonly do this!

... these atoms have unfilled "d" orbitals that may participate in bonding!

- All noble gas compounds (example: XENON compounds with oxygen and fluorine) exhibit this behavior!


EXAMPLES:




- The central SULFUR atom has a share in TWELVE total electrons, not eight!
- The SHAPE of the sulfur hexafluoride molecule in three dimensions agrees with the picture of six fluorine atoms each sharing a pair of electrons with a sulfur center.



$$\begin{array}{r} \text{P: } 5 \\ \text{Cl: } \frac{7 \times 3 = 21}{26} \end{array}$$



This structure obeys the octet rule.



This molecule does NOT obey the octet rule. Phosphorus ends up with ten electrons instead of eight.