MOLAR CONCENTRATION *

- unit: MOLARITY (M): moles of dissolved substance per LITER of solution

∠dissolved substance

$$M = \text{molarity} = \frac{\text{moles of SOLUTE}}{\text{L SOLUTION}}$$

If you have 0.250 L (250 mL) of 6.0 M HCI, how many moles of HCI do you have? 6.0 M HC| \longrightarrow 6.0 ms| HC| = L

If you need 0.657 moles of hydrochloric acid, how many liters of 0.0555 M HCl do you need to measure out?

0.0555 MHCI
$$\rightarrow$$
 0.0555 Mal HCI = L

0.657 mal HC| \times $\frac{L}{0.0555 \text{ mal HCI}} = \frac{11.8 L}{(11800 \text{ mL})}$ This volume's a bit large for lab work!

$$0.657 \text{ mol HCI} \times \frac{L}{6.00 \text{ mol HCI}} = \frac{0.110 L}{(11\overline{0} \text{ mL})}$$


Example: How would we prepare 500. mL of 0.500 M sodium sulfate in water?

Naz Soy: 142.05 g/mol

H20

Dissolve the appropriate amount of sodium sulfate into enough water to make 500. mL of

solution.



A VOLUMETRIC FLASK is a flask that is designed to precisely contain a certain volume of liquid.

VOLUMETRIC FLASKS are used to prepare solutions.

volumetric flask

We will use the VOLUME and the MOLARITY to find out how many MOLES of sodium sulfate need to be dissolved. Then, we can convert MOLES to MASS using FORMULA WEIGHT.

0.500 mol Naz
$$504 = L$$
 mL = $10^{-3}L$
 $500.mL \times \frac{10^{-3}L}{mL} \times \frac{0.500 \, mol \, Naz \, 504}{L} = 0.250 \, mol \, Naz \, 504$

$$\frac{142.05 \, g \, Na_2 50 \, q = mol \, Na_2 50 \, q}{0.250 \, mol \, Na_2 50 \, q \, x \frac{142.05 \, g \, Na_2 50 \, q}{mol \, Na_2 50 \, q} = 35.5 \, g \, No_2 50 \, q}$$

Weigh 35.5 g sodium sulfate into 500 mL volumetric flask, then add water to the mark.

More on MOLARITY

To prepare a solution of a given molarity, you generally have two options:

- Weigh out the appropriate amount of solute, then dilute to the desired volume with solvent (usually water)
- Take a previously prepared solution of known concentration and DILUTE it with solvent to form a new solution

- Use DILUTION EQUATION

The dilution equation is easy to derive with simple algebra.

... but when you dilute a solution, the number of moles of solute REMAINS CONSTANT. (After all, you're adding only SOLVENT)

$$M_1 V_1 = M_2 V_2$$
 Since the number of moles of solute stays before after the same, this equality must be true!

$$M_1 V_1 = M_2 V_2$$
 ... the "DILUTION EQUATION"

M, = molarity of concentrated solution

 $\sqrt{}$ volume of concentrated solution

M 2 = molarity of dilute solution

V2 = volume of dilute solution (total valume, not volume at added solvent!)

The volumes don't HAVE to be in liters, as long as you use the same volume UNIT for both volumes!

Example: Take the 0.500 M sodium sulfate we discussed in the previous example and dilute it to make 150. mL of 0.333 M solution. How many mL of the original solution will we need to dilute?

$$M_1V_1 = M_2V_2$$
 $(0.500M)V_1 = (0.3333M)(150.mL)$
 $V_1 = ?$
 $V_2 = 150.mL$
 $V_1 = 99.9mL$ of 0.500M Naz504

Measure out 99.9 mL of 0.500 M sodium sulfate, then add enough water to get a total volume of 150. mL (could do this in a graduated cylinder!)

- Chemical reactions proceed on an ATOMIC basis, NOT a mass basis!
- To calculate with chemical reactions (i.e. use chemical equations), we need everything in terms of ATOMS ... which means MOLES of atoms

2 Al (s) +3 Br₂(1)
$$\rightarrow$$
 2 Al Br₃(s)
Toefficients are in terms of atoms and molecules!
2 atoms Al = 3 molecules Br₂ = 2 formula units Al Br₃
2 mol Al = 3 mol Br₂ = 2 mol Al Br₃

- To do chemical calculations, we need to:
 - Relate the amount of substance we know (mass or volume) to a number of moles
 - Relate the moles of one substance to the moles of another using the equation
 - Convert the moles of the new substance to mass or volume as desired

$$2A(ls) + 3Br_2(l) \longrightarrow 2A(Br_3(s))$$

- * Given that we have 25.0 g of liquid bromine, how many grams of aluminum would we need to react away all of the bromine?
 - Convert grams of bromine to moles: Need formula weight B_{12} , 2×79.90 159.80 $25.09 BC2 \times \frac{mol BC2}{159.80} = 0.15645 \text{ mol BC2}$
 - Use the chemical equation to relate moles of bromine to moles of aluminum $2 \text{ mol } A = 3 \text{ mol } B_2$

Convert moles aluminum to mass: Need formula weight A1:26.98 26.989A1= mol A1

You can combine all three steps on one line if you like!

Things we can do:

If we have	and we need	Use
MASS	MOLES	FORMULA WEIGHT
SOLUTION VOLUME	MOLES	MOLAR CONCETRATION (MOLARITY)
MOLES OF A	MOLES OF B	BALANCED CHEMICAL EQUATION

101 Example:

How many milliliters of 6.00M hydrochloric acid is needed to completely react with 25.0 g of sodium carbonate?

$$=$$
 2 H(1(aq)+ Na₂(O₃(s) \longrightarrow H₂O(l) + (O₂(g)+ 2NaCl(aq))

- 1 Convert 25.0 g sodium carbonate to moles. Use FORMULA WEIGHT.
- 2 Convert moles sodium carbonate to moles HCI. Use CHEMICAL EQUATION.
- 3 Convert moles HCI to volume solution. Use MOLARITY (6.00 M)

$$\begin{array}{c|c}
\hline
 Na_{2}(0_{3}; N_{4}-2x2z.99 \\
 (-1x|2.0| \\
 0-3x|6.00 \\
\hline
 105.99_{g}N_{62}(0_{3}=mol\ N_{62}(0_{3}=mol\ N_$$