22.4 grams of benzene are reacted with excess nitric acid. If 31.6 grams of nitrobenzene are collected from the reaction, what is the percent yield?

We need to calculate the THEORETICAL YIELD of nitrobenzene, starting with 22.4 grams of benzene (the starting material)

$$\frac{1}{3} | 123.1 | 19 | (6 H_6 = mol (6 H_6 = mol (6 H_6 = mol (6 H_6 = mol (6 H_6 NO_2))) | (6 H_6 NO_2 = mol (6 H_6 NO_2) | (6 H_6 NO_2) | (123.1 | 19 ($$

Now calculate the percent yield:

- electrolytes: substances that dissolve in water to form charge-carrying solutions
- * Electrolytes form ions in solution (ions that are mobile are able to carry charge!). These IONS can interact with one another and undergo certain kinds of chemistry!

IONIC THEORY

- the idea that certain compounds DISSOCIATE in water to form free IONS

Strong vs weak?

- If an electrolyte COMPLETELY IONIZES in water, it's said to be STRONG
- If an electrolyte only PARTIALLY IONIZES in water, it's said to be WEAK
- Both kinds of electrolyte undergo similar kinds of chemistry.

What kinds of compounds are electrolytes?

MOLECULAR COMPOUNDS

- Most molecular compounds are NONELECTROLYTES they don't ionize in water
- -ACIDS and BASES will ionize in water. Most of these are WEAK ELECTROLYTES, but there are a few STRONG ACIDS and STRONG BASES.


$$\begin{array}{c} (12 \text{ Hz2})_{11}(s) \xrightarrow{\text{Hz0}} (12 \text{ Hz2})_{11}(aq) & \text{... nonelectroiyte} \\ H(2 \text{ Hz})_{2}(l) \xrightarrow{\text{Hz0}} H^{+}(aq) + (2 \text{ Hz0})^{-}(aq) \end{array}$$

... acetic acid (electrolyte)


IONIC COMPOUNDS

- SOLUBLE ionic compounds are STRONG ELECTROLYTES they completely ionize in water.
- Not all ionic compounds are water soluble, however!

- provides an easy-to-understand MECHANISM for certain kinds of chemical reactions.
 - "Exchange" reactions. (a.k.a "double replacement" reactions)



Looking a bit more closely...



(The net ionic equation shows only ions and substances that change during the course of the reaction!)

- The net ionic equation tells us that any source of aqueous silver and chloride ions will exhibit this same chemistry, not just silver nitrate and sodium chloride!

- molecular equations: Represent all substances (even ionic substances) as if they were molecules. Include spectator ions, and do not show charges on ions. Traditional chemical equations.
- ionic equations: Show all free ions including spectators in a chemical reaction. Molecules and WEAK electrolytes are shown as molecules. STRONG electrolytes (like HCl) are shown as ions. Ions that are part of undissolved ionic compounds are shown as molecules.
- NET ionic equation: An ionic equation that leaves out spectator ions. Intended to show only things that actually change in a reaction.

Ag NO₃ (aq) + NaCl (aq)
$$\rightarrow$$
 Ag(l(s) + NaNo₃ (aq)

Ag (aq) + No₃ (aq) + Na^t (aq) + Cl (aq) \rightarrow Ag(l(s) + Na^t (aq) + No₃ (aq)

Ag (aq) + Cl (aq) \rightarrow Ag(l(s)

* You can get from the complete ionic equation to the net ionic equation by crossing out the spectator ions on both sides.

How can I tell if an ionic compound dissolves in water?

consult experimental data: "solubility rules"!

A few of the "rules"...

- Compounds that contain a Group IA cation (or ammonium) are soluble
- Nitrates and acetates are soluble
- Carbonates, phosphates, and hydroxides tend to be insoluble

... or see the web site for a solubility chart.

#8 - hydroxides generally insoluble, except Group IA, ammonium, calcium strontium, barium

Conclusion: iron(III) hydroxide is insoluble.

#3 - lodides usually dissolve, exceptions are silver, mercury, lead

Conclusion: silver(I) iodide is INSOLUBLE

#2 - acetates are soluble, no common exceptions.

Conclusion: calcium acetate is soluble.

#5 - Most carbonates are insoluble

Conclusion - barium carbonate is insoluble.

Exchange Chemistry

- Three kinds of exchange chemistry.

(I) PRECIPITATION Formation of IONIC SOLID

(2) ACID/BASE or NEUTRALIZATION

Formation of MOLECULES!

GAS FORMATION (formation of unstable molecules)

SOME (but not all) reactions that form gases
are examples of exchange chemistry.

Just because you mix together two ionic compounds does NOT mean that a reaction will occur. You need a DRIVING FORCE for a reaction.


PRECIPITATION REACTIONS

- driving force is the formation of an insoluble ionic compound.

3Mg(
$$1_2(aq) + 2Na_3 PO_4(aq) \rightarrow 6Na(|aq) + Mg_3(PO_4)_2(S)$$
ions: Mg C1 Na PO43 This reaction is driven by the formation of INSOLUBLE magnesium phosphate.

When you're trying to complete a precipitation reaction:

- (1) Write the IONS that form when the reactants are dissolved.
- Make NEW compounds by pairing up cations with anions. Don't forget that the positive and negative charges must balance each other out!
- (3) Use the solubility rules to determine the PHASE of each new compound solid or aqueous.
- (4) Balance the overall equation.



So, no solid forms here. All possible combinations of these four ions result in compounds that dissolve readily in water.

$$m_g^{2+}Cl^{-}$$
 + N_a^{+} $C_2^{+}N_3^{0}$ N_a^{+} M_a^{+} M_a^{+}


★ We will learn about other driving forces than the formation of solid, but these driving forces do not apply to this reaction

ACID/BASE REACTIONS (also called NEUTRALIZATION REACTIONS)

- There are several stable molecules that may be formed in double replacement reactions, but the most common is WATER!
- Double replacement reactions that form water are also called "neutralizations"

HA + BOH
$$\longrightarrow$$
 H2O + BA acid base salt in hoh" ionic compound

* To make water (H_2O), you need a source of hydrogen ion (H^4) and hydroxide ion (OH^5)



ACIDS

- compounds that release hydrogen ion (H), when dissolved in water.

Properties of acids:

- Corrosive: React with most metals to give off hydrogen gas
- Cause chemical burns on contact
- Taste sour (like citrus citric acid!)
- Changes litmus indicator to RED

BASES

- Substances that release hydroxide ion (OH*) when dissolved in water

Properties of bases:

- Caustic: Attack and dissolve organic matter (think lye, which is NaOH)
- Cause skin/eye damage on contact
- Taste bitter
- changes litmus indicator to BLUE

Due to the dissolving action of base on your skin, bases will feel "slippery". The base ITSELF is not particularly slippery, but what's left of your skin IS!

ACID/BASE or NEUTRALIZATION reactions continued

- the driving force of these reactions is the formation of water molecules.

$$H^{+}(aq) + OH^{-}(aq) \longrightarrow H_{2}O(Q)$$
 Net ionic equation
From the acid From the base

H2S0y(ay) + 2NaOH(ay)
$$\longrightarrow$$
 2H2O(l) + Na2SOy(aq) Vions: H+ SOy2- Na+ OH-

- How can this reaction be detected?
 - pH detector (indicator paper, etc.)
 - do the products have similar chemical properties to the reactants?
 - release of heat!

... formation of water is usually accompanied by a release of heat

GAS FORMATION / OTHER MOLECULES

- There are a few other molecules that can be made with exchange-type chemistry.
- Most of these molecules are unstable and can break apart to form gases.
- Formation of a weak acid:
 - The formation of ANY weak acid in an exchange-type reaction can be a driving force.
 - Some weak acids are unstable and can break apart into gas molecules.

$$H_2(o_3 Lag) \longrightarrow H_2(l) + Co_2(g)$$
 Gas bubbles can leave solution!

... but how would you form carbonic acid in an exchange-type reaction?

$$H_2SO_4(a_4)+2NaH(O_3(a_4)) \rightarrow Na_2SO_4(a_4)+2H_2CO_3(a_4)$$
 $H^+SO_4^2-Na^+H(O_3^-)$

... but when we mix sulfuric acid and sodium bicarbonate, we observe BUBBLES. We need to write an equation that agrees with our observations. We know that carbonic acid decomposes, so we go ahead and put that into our equation.


$$H_2(o_3lag) \longrightarrow H_2O(l) + (o_2(g))$$
 $H_2So_4(ag) + 2NaH(o_3lag) \rightarrow Na_2So_4(ag) + 2H_2O(l) + 2(o_2(g))$

Other molecules of interest:

$$H_2$$
 SO $_3$: sulfurous acid - React an ACID with a SULFITE

$$H_2So_3(ag) \rightarrow H_2O(\ell) + So_2(g)$$

 H_2S hydrogen sulfide (gas) - React an ACID with a SULFIDE



This reaction is driven by the formation of the weak acid (molecule) CARBONIC ACID, and its decomposition into WATER and CARBON DIOXIDE GAS.