Calculations with measurements

When you calculate something using measured numbers, you should try to make sure the ANSWER reflects the quality of the data used to make the calculation.

An ANSWER is only as good as the POOREST measurement that went into finding that answer!

$$\begin{array}{r}
14.206 \pm 0.001 \\
154.72 \pm 0.01 \\
1.6 \pm 0.1 \\
\pm 0.222 \pm 0.001 \\
170.748
\end{array}$$

How should we report this answer? How much uncertainty is in this answer?

✓ If you add an uncertain number to either a certain or an uncertain number, then the result is uncertain!

 \star If you add certain numbers together, the result is certain!

For addition and subtraction, round FINAL ANSWERS to the same number of decimal places as the measurement with the fewest decimal places. This will give an answer that indicates the proper amount of uncertainty.

For multiplication and division, round FINAL ANSWERS to the same number of SIGNIFICANT FIGURES as the measurement with the fewest SIGNIFICANT FIGURES!

$$4 \quad 3 \quad 3 \\ 15.62 \times 0.0667 \times 35.0 = 36.46489$$

How should we report this answer?
$$36.5$$

$$25.4 \times 0.00023 \times 15.201 = 0.088804242$$

How should we report this answer?
$$0.089$$

Remember that lead zeros can't be significant. So the first sig fig is the leftmost "8". Neither zero on the left is significant.

27

A few more math with significant figures examples:

$$\frac{5}{15047} \times \frac{9}{11} \times 0.9876 = 163464.5892$$

$$\frac{160000}{OR} \frac{16000}{OR} \frac{16000}{PR} \frac{16000}{PR} \frac{16000}{PR} \frac{160000}{PR} \frac{16000}{PR} \frac{160000}{PR} \frac{160000}{PR} \frac{160000}{PR} \frac{16000}{PR} \frac{16$$

Exact Numbers

- Some numbers do not have any uncertainty. In other words, they weren't measured!

1) Numbers that were determined by COUNTING!

2) Numbers that arise from DEFINITIONS, often involving relationships between units 12 in = 1 FE Km = 10³m All metric prefixes are exact!

V exactly 4

How many blocks are to the left?

³⁰ Example

You'll need to round the answer to the right number of significant figures! Convert 4.45 m to in, assuming that 2.54 cm = 1 in

2.54 cm zin
$$Cm = 10^{-2}m$$

4.45 m x $\frac{Cm}{10^{-2}m}$ x $\frac{in}{2.54}$ gm = 175.1968504 in
 $\int_{3}^{1} \int_{-\infty}^{0} \int_{-\infty}^{\infty} \int_{-\infty$

Usually, in unit conversions the answer will have the same number of significant figures as the original measurement did.

EXCEPTION: Temperature conversions, since these often involve ADDTION (different rule!)

A note on rounding: If possible, try to round only at the END of a multiple-step calculation. Avoid rounding intermediate numbers if possible, since extra rounding introduces RANDOM ERROR into the calculation.

- 1808: Publication of Dalton's "A New System of Chemical Philosophy", which contained the atomic theory

- Dalton's theory attempted to explain two things:



CONSERVATION OF MASS

- The total amount of mass remains constant in any process, chemical or physical!



<u>LAW OF DEFINITE PROPORTIONS (also called the LAW OF CONSTANT</u> COMPOSITION): All pure samples of a given compound contain the same proportion of elements by mass The parts of Dalton's theory

() Matter is composed of small, chemically indivisible <u>ATOMS</u>

Description ELEMENTS are kinds of matter that contain only a single kind of atom. All the atoms of an element have identical chemical properties.

3 COMPOUNDS are kinds of matter that are composed of atoms of two or more ELEMENTS which are combined in simple, whole number ratios.

Most importantly,

CHEMICAL REACTIONS are REARRANGEMENTS of atoms to form new compounds.

- Atoms are not gained or lost during a chemical reaction.
- Atoms do not change their identity during a chemical reaction.
- All the atoms that go into a chemical reaction must go out again!